



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 48/00, C12N 15/12, 15/31, 15/33, 15/48		A1	(11) International Publication Number: WO 94/16737
			(43) International Publication Date: 4 August 1994 (04.08.94)
(21) International Application Number: PCT/US94/00899		(71)(72) Applicants and Inventors: WEINER, David, B. [US/US]; 717 Beacon Lane, Merion, PA 19066 (US). WILLIAMS, William, V. [US/US]; 25 Sycamore Road, Havertown, PA 19083 (US). WANG, Bin [CN/US]; 610 Covington Road, Havertown, PA 19083 (US). CONEY, Leslie, R. [US/US]; 1000 Conestoga Road, Rosemont, PA 19101 (US). MERVA, Michael, J. [US/US]; 5430 Hermit Terrace, Apartment 1F, Philadelphia, PA 19128 (US). ZURAWSKI, Vincent, R., Jr. [US/US]; 728 Northgate Road, Westtown, PA 19395 (US).	
(22) International Filing Date: 26 January 1994 (26.01.94)		(74) Agents: ELDERKIN, Dianne, B. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris, One Liberty Place - 46th floor, Philadelphia, PA 19103 (US).	
(30) Priority Data: 08/008,342 26 January 1993 (26.01.93) US 08/029,336 11 March 1993 (11.03.93) US 08/093,235 15 July 1993 (15.07.93) US 08/124,962 21 September 1993 (21.09.93) US 08/125,012 21 September 1993 (21.09.93) US			
(60) Parent Applications or Grants (63) Related by Continuation US 08/008,342 (CIP) Filed on 26 January 1993 (26.01.93) US 08/029,336 (CIP) Filed on 11 March 1993 (11.03.93) US 08/093,235 (CIP) Filed on 15 July 1993 (15.07.93) US 08/124,962 (CIP) Filed on 21 September 1993 (21.09.93) US 08/125,012 (CIP) Filed on 21 September 1993 (21.09.93)		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
		Published With international search report.	
(54) Title: COMPOSITIONS AND METHODS FOR DELIVERY OF GENETIC MATERIAL			
(57) Abstract Methods of introducing genetic material into cells of an individual and compositions and kits for practicing the same are disclosed. The methods comprise the steps of contacting cells of an individual with a polynucleotide function enhancer and administering to the cells, a nucleic acid molecule that is free of retroviral particles. The nucleic acid molecule comprises a nucleotide sequence that encodes a protein that comprises at least one epitope that is identical or substantially similar to an epitope of a pathogen antigen or an antigen associated with a hyperproliferative or autoimmune disease, a protein otherwise missing from the individual due to a missing, non-functional or partially functioning gene, or a protein that produces a therapeutic effect on an individual. Methods of prophylactically and therapeutically immunizing an individual against HIV are disclosed. Pharmaceutical compositions and kits for practicing methods of the present invention are disclosed.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	CN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TC	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Compositions and Methods for Delivery of Genetic Material

Field of the Invention

The present invention relates to compositions and methods for introducing genetic material into the cells of an individual. The compositions and methods of the invention can be used to deliver protective and/or therapeutic agents including genetic material that encodes protein targets for immunization and therapeutic proteins.

Background of the Invention

10 The direct introduction of a normal, functional gene into a living animal has been studied as a means for replacing defective genetic information. In some studies, DNA is introduced directly into cells of a living animal without the use of a viral particle or other infectious vector. Nabel, 15 E.G., et al., (1990) *Science* 249:1285-1288, disclose site-specific gene expression *in vivo* of a beta-galactosidase gene that was transferred directly into the arterial wall in mice. Wolfe, J.A. et al., (1990) *Science* 247:1465-1468, disclose expression of various reporter genes that were directly 20 transferred into mouse muscle *in vivo*. Acsadi G., et al., (1991) *Nature* 352:815-818, disclose expression of human dystrophin gene in mice after intramuscular injection of DNA constructs. Wolfe, J.A., et al., 1991 *BioTechniques* 11(4):474-485, which is incorporated herein by reference,

- 2 -

refers to conditions affecting direct gene transfer into rodent muscle *in vivo*. Felgner, P.L. and G. Rhodes, (1991) *Nature* 349:351-352, disclose direct delivery of purified genes *in vivo* as drugs without the use of retr viruses.

5 The use of direct gene transfer as an alternative anti-pathogen vaccination method has been suggested. Use of direct gene transfer by single injection is suggested as a possible vaccination strategy against HIV. A cellular immune response to HIV gp120 resulting from introduction of plasmid
10 DNA encoding the same into cells is reported to have been observed. PCT International Application Number PCT/US90/01515 published October 4, 1990 discloses methods of immunizing an individual against pathogen infection by directly injecting
15 naked polynucleotides into the individual's cells in a single step procedure. The use of transfecting agents other than lipofectins is specifically excluded from the disclosed methods. The stimulation of inoculated cells is neither disclosed nor suggested. An HIV vaccine is disclosed which consists of the introduction of polynucleotides that encode
20 the viral protein gp120. The operability of this vaccine is not evidenced.

 Thomason, D.B. et al., (1990) *Cell Physiol.* 27:C578-581 and PCT Patent Application Serial No. WO 91/12329 disclose administering bupivacaine to muscle cells in order to induce
25 satellite cell proliferation as part of a retroviral-mediated gene delivery protocol.

Summary of the Invention

 The present invention relates to methods of introducing genetic material into the cells of an individual.
30 The methods comprises the steps of contacting cells of said individual with a polynucleotide function enhancer agent, which is preferably an agent that facilitates the uptake of DNA by the cells or enhances an inflammatory response, and administering to the cells, a nucleic acid molecule that
35 comprises a nucleotide sequence that either encodes a desired peptide or protein, or serves as a template for functional nucleic acid molecules. The nucleic acid molecule is administered free from retr viral particles. The desired

- 3 -

protein may either be a protein which functions within the individual or it serves as a target for an immune response.

The present invention relates to a method of immunizing an individual against a pathogen. The method comprises the steps of contacting cells of said individual with a polynucleotide function enhancer agent, which is preferably an agent that facilitates the uptake of DNA by the cells or enhances the immune response, and administering to the cells, a nucleic acid molecule that comprises a nucleotide sequence that encodes a peptide which comprises at least an epitope identical or substantially similar to an epitope displayed on a pathogen antigen and is operatively linked to regulatory sequences. The nucleic acid molecule is capable of being expressed in the cells of the individual.

The present invention relates to a method of immunizing a human against HIV. The method comprises the steps of administering to a human a nucleic acid molecule that comprises a nucleotide sequence that encodes at least one peptide that comprises at least one epitope identical or substantially similar to an epitope displayed on an HIV protein operatively linked to regulatory sequences.

The present invention relates to a method of immunizing a human against HIV. The method comprises the steps of administering two different nucleic acid molecules to different cells of the human. Each nucleic acid molecule comprises a nucleotide sequence that encodes at least one peptide which comprises at least one epitope identical or substantially similar to an epitope displayed on an HIV protein operatively linked to regulatory sequences. The different nucleic acid molecules each comprise different nucleotide sequences that encode at least one different peptide from the other and are each capable of being expressed in human cells.

The present invention relates to methods of immunizing an individual against a hyperproliferative disease or an autoimmune disease. The methods comprise the steps of administering to cells of an individual, a nucleic acid

- 4 -

molecule that comprises a nucleotide sequence that encodes a peptide that comprises at least an epitope identical or substantially similar to an epitope displayed on a hyperproliferative disease-associated protein or an autoimmune disease-associated protein, respectively, and is operatively linked to regulatory sequences; the nucleic acid molecule being capable of being expressed in the cells.

The present invention relates to methods of treating an individual suffering from a disease comprising the steps of contacting cells of said individual with a polynucleotide function enhancer agent, which is preferably an agent that facilitates the uptake of DNA by the cells or enhances an inflammatory response, and administering to cells of an individual, a nucleic acid molecule that comprises a nucleotide sequence which functions in place of a defective gene or which encodes a molecule that produces a therapeutic effect in the individual and is operatively linked to regulatory sequences; the nucleic acid molecule being capable of being expressed in the cells.

The present invention relates to pharmaceutical compositions which comprise a nucleic acid molecule and a polynucleotide function enhancer. The present invention relates to pharmaceutical kits which comprise a container comprising a nucleic acid molecule and a container comprising a polynucleotide function enhancer.

The present invention relates to prophylactic and therapeutic HIV vaccines comprising a pharmaceutically acceptable carrier or diluent and a nucleic acid molecule that encodes one or more peptides that each comprises at least an epitope identical or substantially similar to an epitope displayed on at least one HIV protein operatively linked to regulatory sequences; the nucleic acid molecule being capable of being expressed in human cells.

The present invention relates to prophylactic and therapeutic HIV vaccines comprising two inoculants. The first inoculant comprises a pharmaceutically acceptable carrier or diluent and a first nucleic acid molecule. The first nucleic

- 5 -

acid molecule comprises a nucleotide sequence that encodes one or more peptides that each comprises at least an epitope identical or substantially similar to an epitope displayed on at least one HIV protein operatively linked to regulatory sequences; the nucleic acid molecule being capable of being expressed in human cells. The second inoculant comprises a pharmaceutically acceptable carrier or diluent and a second nucleic acid molecule. The second nucleic acid molecule comprises a nucleotide sequence that encodes one or more peptides that each comprises at least an epitope identical or substantially similar to an epitope displayed on at least one HIV protein operatively linked to regulatory sequences; the nucleic acid molecule being capable of being expressed in human cells. The first and second nucleic acid molecules are different and encode different peptides.

Brief Description of the Drawings

Figure 1A is a diagram depicting the construction of plasmid pM160 which was produced by inserting a PCR-generated fragment that encodes the HIV-HXB2 glycoprotein gp160 into plasmid pMAMneoBlue (Clonetech).

Figure 1B is a photograph of an autoradiogram of a Western blot of whole cell lysates of cells transfected with the pM160 plasmid (3G7 cells) versus vector-alone transfected cells (TE671 cells) showing production of gp120 and gp41 in 3G7 cells and not in TE671 cells.

Figure 2 is a photograph of an autoradiogram showing immunoprecipitations of serum antibodies binding to ^{125}I -gp160.

Figures 3A - 3E are graphs showing ELISA results binding different sera to various proteins immobilized on microtiter plates.

Figures 4A and 4B are photographs of MT-2 cells infected with TCID₅₀HIV-1/III_B cell-free virus that was preincubated with serial dilutions of antisera.

Figure 4C is a graph illustrating the neutralization values (V_n/V_0) versus the dilution factors from results using

- 6 -

control serum (x = pMAMneoBlue vector-immunized mice) and test sera (0 = pM160-immunized mice).

Figures 4D - 4G are photographs of H9/III₂ cells used in experiments to examine syncytial inhibition using sera from immunized and control animals.

Figure 5 is a chart depicting the survival of immunized and non-immunized mice challenged with HIV gp160-labelled and unlabelled tumor cells. Mice were immunized with recombinant gp160 protein, vector DNA only or recombinant vector comprising DNA encoding gp160. SP2/0 tumor cells or SP2/0-gp160 (SP2/0 cells transfected with DNA encoding gp160 and expressing gp160) tumor cells were introduced into the mice.

Figure 6 is a plasmid map of pGAGPOL.rev.

Figure 7 is a plasmid map of pENV.

Figure 8 is shows four backbones, A, B, C and D, used to prepare genetic construct.

Figure 9 shows four inserts, 1, 2, 3 and 4 which are inserted into backbones to produce genetic constructs.

20 Detailed Description of the Invention

The present invention relates to a method of introducing nucleic acid molecules into the cells of an animal which provides for the high level of uptake and function of the nucleic acid molecules. The method of the present invention comprises the steps of administering nucleic acid molecules that are free from viral particles, particularly retroviral particles, to the cell of an individual in conjunction with administration of a co-agent which enhances the inflammatory response and/or enhances expression of the nucleic acid molecule in the tissue and/or facilitates the uptake of the nucleic acid molecule by the cell. Preferred embodiments of the present invention provide methods of delivering nucleic acid molecules to cells of an individual without the use of infectious agents.

35 Nucleic acid molecules which are delivered to cells according to the invention may serve as: 1) genetic templates

- 7 -

for proteins that function as prophylactic and/or therapeutic immunizing agents; 2) replacement copies of defective, missing or non-functioning genes; 3) genetic templates for therapeutic proteins; 4) genetic templates for antisense molecules; or 5) genetic templates for ribozymes. In the case of nucleic acid molecules which encode proteins, the nucleic acid molecules preferably comprise the necessary regulatory sequences for transcription and translation in the cells of the animal. In the case of nucleic acid molecules which serve as templates for antisense molecules and ribozymes, such nucleic acid molecules are preferably linked to regulatory elements necessary for production of sufficient copies of the antisense and ribozyme molecules encoded thereby respectively. The nucleic acid molecules are free from retroviral particles and preferably provided as DNA in the form of plasmids.

The co-agent is also referred to herein as a "polynucleotide function enhancer" or "PFE". A PFE is a compound or composition which enhances the inflammatory response and/or enhances expression of the nucleic acid molecule in the tissue and/or facilitates the uptake of the nucleic acid molecule by the cell and preferably has more than one of these properties. Polynucleotide function enhancers that facilitate DNA and RNA uptake by cells and stimulate cell division and replication are also referred to as cell stimulating agents. Preferred co-agents according to the present invention are selected from the group consisting of benzoic acid esters and anilides. In preferred embodiments, the PFE is bupivacaine.

According to some aspects of the present invention, compositions and methods are provided which prophylactically and/or therapeutically immunize an individual against a pathogen or abnormal, disease-related cell. The genetic material encodes a peptide or protein that shares at least an epitope with an immunogenic protein found on the pathogen or cells to be targeted. The genetic material is expressed by the individual's cells and serves as an immunogenic target against which an immune response is elicited. The resulting

- 8 -

immune response is broad based: in addition to a humoral immune response, both arms of the cellular immune response are elicited. The methods of the present invention are useful for conferring prophylactic and therapeutic immunity. Thus, a method of immunizing includes both methods of protecting an individual from pathogen challenge, or occurrence or proliferation of specific cells as well as methods of treating an individual suffering from pathogen infection, hyperproliferative disease or autoimmune disease.

The present invention is useful to elicit broad immune responses against a target protein, i.e. proteins specifically associated with pathogens or the individual's own "abnormal" cells. The present invention is useful to immunize individuals against pathogenic agents and organisms such that an immune response against a pathogen protein provides protective immunity against the pathogen. The present invention is useful to combat hyperproliferative diseases and disorders such as cancer by eliciting an immune response against a target protein that is specifically associated with the hyperproliferative cells. The present invention is useful to combat autoimmune diseases and disorders by eliciting an immune response against a target protein that is specifically associated with cells involved in the autoimmune condition.

Some aspects of the present invention relate to gene therapy; that is, to compositions for and methods of introducing nucleic acid molecules into the cells of an individual exogenous copies of genes which either correspond to defective, missing, non-functioning or partially functioning genes in the individual or which encode therapeutic proteins, i.e. proteins whose presence in the individual will eliminate a deficiency in the individual and/or whose presence will provide a therapeutic effect on the individual thereby providing a means of delivering the protein by an alternative means from protein administration.

As used herein the term "desired protein" is meant to refer to peptides and protein encoded by gene constructs of the present invention which either act as target proteins

- 9 -

for an immune response or as a therapeutic or compensating protein in gene therapy regimens.

According to the present invention, DNA or RNA that encodes a desired protein is introduced into the cells of an individual where it is expressed, thus producing the desired protein. The DNA or RNA encoding the desired protein is linked to regulatory elements necessary for expression in the cells of the individual. Regulatory elements for DNA expression include a promoter and a polyadenylation signal. In addition, other elements, such as a Kozak region, may also be included in the genetic construct.

As used herein, the term "genetic construct" refers to the DNA or RNA molecule that comprises a nucleotide sequence which encodes the desired protein and which includes initiation and termination signals operably linked to regulatory elements including a promoter and polyadenylation signal capable of directing expression in the cells of the vaccinated individual.

As used herein, the term "expressible form" refers to gene constructs which contain the necessary regulatory elements operable linked to a coding sequence that encodes a target protein, such that when present in the cell of the individual, the coding sequence will be expressed.

As used herein, the term "genetic vaccine" refers to a pharmaceutical preparation that comprises a genetic construct that comprises a nucleotide sequence that encodes a target protein including pharmaceutical preparations useful to invoke a therapeutic immune response.

As used herein, the term "genetic therapeutic" refers to a pharmaceutical preparation that comprises a genetic construct that comprises a nucleotide sequence that encodes a therapeutic or compensating protein.

As used herein, the term "target protein" refers to a protein against which an immune response can be elicited. The target protein is an immunogenic protein which shares at least an epitope with a protein from the pathogen or undesirable cell-type such as a cancer cell or a cell involved

- 10 -

in autoimmune disease against which immunization is required. The immune response directed against the target protein will protect the individual against and treat the individual for the specific infection or disease with which the target protein is associated.

As used herein, the term "sharing an epitope" refers to proteins which comprise at least one epitope that is identical to or substantially similar to an epitope of another protein.

As used herein, the term "substantially similar epitope" is meant to refer to an epitope that has a structure which is not identical to an epitope of a protein but nonetheless invokes an cellular or humoral immune response which cross reacts to that protein.

As used herein, the term "therapeutic protein" is meant to refer to proteins whose presence confers a therapeutic benefit to the individual.

As used herein, the term "compensating protein" is meant to refer to proteins whose presence compensates for the absence of a fully functioning endogenously produced protein due to an absent, defective, non-functioning or partially functioning endogenous gene.

Genetic constructs comprise a nucleotide sequence that encodes a desired protein operably linked to regulatory elements needed for gene expression. Accordingly, incorporation of the DNA or RNA molecule into a living cell results in the expression of the DNA or RNA encoding the desired protein and thus, production of the desired protein.

When taken up by a cell, the genetic construct which includes the nucleotide sequence encoding the desired protein operably linked to the regulatory elements may remain present in the cell as a functioning extrachromosomal molecule or it may integrate into the cell's chromosomal DNA. DNA may be introduced into cells where it remains as separate genetic material in the form of a plasmid. Alternatively, linear DNA which can integrate into the chromosome may be introduced into the cell. When introducing DNA into the cell, reagents which

- 11 -

promote DNA integration into chromosomes may be added. DNA sequences which are useful to promote integration may also be included in the DNA molecule. Alternatively, RNA may be administered to the cell. It is also contemplated to provide
5 the genetic construct as a linear minichromosome including a centromere, telomeres and an origin of replication.

The molecule that encodes a desired protein may be DNA or RNA which comprise a nucleotide sequence that encodes the desired protein. These molecules may be cDNA, genomic
10 DNA, synthesized DNA or a hybrid thereof or an RNA molecule such as mRNA. Accordingly, as used herein, the terms "DNA construct", "genetic construct" and "nucleotide sequence" are meant to refer to both DNA and RNA molecules.

The regulatory elements necessary for gene
15 expression of a DNA molecule include: a promoter, an initiation codon, a stop codon, and a polyadenylation signal. In addition, enhancers are often required for gene expression. It is necessary that these elements be operably linked to the sequence that encodes the desired proteins and that the
20 regulatory elements are operably in the individual to whom they are administered.

Initiation codons and stop codon are generally considered to be part of a nucleotide sequence that encodes the desired protein. However, it is necessary that these
25 elements are functional in the individual to whom the gene construct is administered. The initiation and termination codons must be in frame with the coding sequence.

Promoters and polyadenylation signals used must be functional within the cells of the individual.

30 Examples of promoters useful to practice the present invention, especially in the production of a genetic vaccine for humans, include but are not limited to promoters from Simian Virus 40 (SV40), Mouse Mammary Tumor Virus (MMTV) promoter, Human Immunodeficiency Virus (HIV) such as the HIV
35 Long Terminal Repeat (LTR) promoter, Moloney virus, ALV, Cytomegalovirus (CMV) such as the CMV immediate early promoter, Epstein Barr Virus (EBV), Rous Sarcoma Virus (RSV)

- 12 -

as well as promoters from human genes such as human Actin, human Myosin, human Hemoglobin, human muscle creatine and human metallothionein.

Examples of polyadenylation signals useful to practice the present invention, especially in the production of a genetic vaccine for humans, include but are not limited to SV40 polyadenylation signals and LTR polyadenylation signals. In particular, the SV40 polyadenylation signal which is in pCEP4 plasmid (Invitrogen, San Diego CA), referred to as the SV40 polyadenylation signal, is used.

In addition to the regulatory elements required for DNA expression, other elements may also be included in the DNA molecule. Such additional elements include enhancers. The enhancer may be selected from the group including but not limited to: human Actin, human Myosin, human Hemoglobin, human muscle creatine and viral enhancers such as those from CMV, RSV and EBV.

Genetic constructs can be provided with mammalian origin of replication in order to maintain the construct extrachromosomally and produce multiple copies of the construct in the cell. Plasmids pCEP4 and pREP4 from Invitrogen (San Diego, CA) contain the Epstein Barr virus origin of replication and nuclear antigen EBNA-1 coding region which produces high copy episomal replication without integration.

In some preferred embodiments, the vector used is selected from those described in Example 46. In aspects of the invention relating to gene therapy, constructs with origins of replication including the necessary antigen for activation are preferred.

In some preferred embodiments related to immunization applications, the genetic construct contains nucleotide sequences that encode a target protein and further include genes for proteins which enhance the immune response against such target proteins. Examples of such genes are those which encode cytokines and lymphokines such as α -interferon, gamma-interferon, platelet derived growth factor

- 13 -

(PDGF), GC-SF, GM-CSF, TNF, epidermal growth factor (EGF), IL-1, IL-2, IL-4, IL-6, IL-8, IL-10 and IL-12. In some embodiments, it is preferred that the gene for GM-CSF is included in genetic constructs used in immunizing compositions.

An additional element may be added which serves as a target for cell destruction if it is desirable to eliminate cells receiving the genetic construct for any reason. A herpes thymidine kinase (tk) gene in an expressible form can be included in the genetic construct. The drug gangcyclovir can be administered to the individual and that drug will cause the selective killing of any cell producing tk, thus, providing the means for the selective destruction of cells with the genetic construct.

In order to maximize protein production, regulatory sequences may be selected which are well suited for gene expression in the cells the construct is administered into. Moreover, codons may be selected which are most efficiently transcribed in the cell. One having ordinary skill in the art can produce DNA constructs which are functional in the cells.

In order to test expression, genetic constructs can be tested for expression levels *in vitro* using tissue culture of cells of the same type as those to be administered. For example, if the genetic vaccine is to be administered into human muscle cells, muscle cells grown in culture such as solid muscle tumors cells of rhabdomyosarcoma may be used as an *in vitro* model to measure expression level.

The genetic constructs used in the present invention are not incorporated within retroviral particles. The genetic constructs are taken up by the cell without retroviral particle-mediated insertion such as that which occurs when retrovirus particles with retroviral RNA that is incorporated in retroviral particles infects a cell. As used herein, the term "free from retroviral particles" is meant to refer to genetic constructs that are not incorporated within retroviral particles. As used herein, "dissociated from an infectious agent" is meant to refer to genetic material which is not part

- 14 -

of a viral, bacterial or eukaryotic vector, either active, inactivated, living or dead, that is capable of infecting a cell.

In some embodiments, the genetic constructs
5 constitute less than a complete, replicatable viral genome such that upon introduction into the cell, the genetic construct possesses insufficient genetic information to direct production of infectious viral particles. As used herein, the term "incomplete viral genome" is meant to refer to a genetic
10 construct which contains less than a complete genome such that incorporation of such a genetic construct into a cell does not constitute introduction of sufficient genetic information for the production of infectious virus.

In some embodiments, an attenuated viral vaccine may
15 be delivered as a genetic construct which contains enough genetic material to allow for production of viral particles. Delivery of the attenuated vaccine as a genetic construct allows for an easier way to produce large quantities of safe, pure active immunizing product.

20 The genetic construct may be administered with or without the use microprojectiles. It is preferred that the genetic constructs of the present invention may be delivered to the cells of an individual free of solid particles. As used herein, the phrase "free of solid particles" is meant to
25 refer to a liquid that does not contain any solid microprojectile used as a means to perforate, puncture or otherwise pierce the cell membrane of a cell in order to create a port of entry for genetic material into the cell.

The present invention may be used to immunize an
30 individual against all pathogens such as viruses, prokaryote and pathogenic eukaryotic organisms such as unicellular pathogenic organisms and multicellular parasites. The present invention is particularly useful to immunize an individual against those pathogens which infect cells and which are not
35 encapsulated such as viruses, and prokaryote such as gonorrhoea, listeria and shigella. In addition, the present invention is also useful to immunize an individual against

- 15 -

protoz an pathogens which include a stage in the life cycle where they are intracellular pathogens. As used herein, the term "intracellular pathogen" is meant to refer to a virus or pathogenic organism that, at least part of its reproductive or life cycle, exists within a host cell and therein produces or causes to be produced, pathogen proteins. Table 1 provides a listing of some of the viral families and genera for which vaccines according to the present invention can be made. DNA constructs that comprise DNA sequences which encode the peptides that comprise at least an epitope identical or substantially similar to an epitope displayed on a pathogen antigen such as those antigens listed on the tables are useful in vaccines. Moreover, the present invention is also useful to immunize an individual against other pathogens including prokaryotic and eukaryotic protozoan pathogens as well as multicellular parasites such as those listed on Table 2.

In order to produce a genetic vaccine to protect against pathogen infection, genetic material which encodes immunogenic proteins against which a protective immune response can be mounted must be included in the genetic construct. Whether the pathogen infects intracellularly, for which the present invention is particularly useful, or extracellularly, it is unlikely that all pathogen antigens will elicit a protective response. Because DNA and RNA are both relatively small and can be produced relatively easily, the present invention provides the additional advantage of allowing for vaccination with multiple pathogen antigens. The genetic construct used in the genetic vaccine can include genetic material which encodes many pathogen antigens. For example, several viral genes may be included in a single construct thereby providing multiple targets. In addition, multiple inoculants which can be delivered to different cells in an individual can be prepared to collectively include, in some cases, a complete or, more preferably, an incomplete such as a near complete set of genes in the vaccine. For example, a complete set of viral genes may be administered using two constructs which each contain a different half of the genome

- 16 -

which are administered at different sites. Thus, an immune response may be invoked against each antigen without the risk of an infectious virus being assembled. This allows for the introduction of more than a single antigen target and can
5 eliminate the requirement that protective antigens be identified.

The ease of handling and inexpensive nature of DNA and RNA further allow for more efficient means of screening for protective antigens. Genes can be sorted and
10 systematically tested much more easily than proteins. The pathogenic agents and organism for which the vaccine is being produced to protect against is selected and an immunogenic protein is identified. Tables 1 and 2 include lists of some of the pathogenic agents and organisms for which genetic
15 vaccines can be prepared to protect an individual from infection by them. In some preferred embodiments, the methods of immunizing an individual against a pathogen are directed against HIV, HTLV or HBV.

Another aspect of the present invention provides a
20 method of conferring a broad based protective immune response against hyperproliferating cells that are characteristic in hyperproliferative diseases and to a method of treating individuals suffering from hyperproliferative diseases. As used herein, the term "hyperproliferative diseases" is meant
25 to refer to those diseases and disorders characterized by hyperproliferation of cells. Examples of hyperproliferative diseases include all forms of cancer and psoriasis.

It has been discovered that introduction of a genetic construct that includes a nucleotide sequence which
30 encodes an immunogenic "hyperproliferating cell"-associated protein into the cells of an individual results in the production of those proteins in the vaccinated cells of an individual. As used herein, the term "hyperproliferative-associated protein" is meant to refer to proteins that are
35 associated with a hyperproliferative disease. To immunize against hyperproliferative diseases, a genetic construct that includes a nucleotide sequence which encodes a protein that

- 17 -

is associated with a hyperproliferative disease is administered to an individual.

In order for the hyperproliferative-associated protein to be an effective immunogenic target, it must be a
5 protein that is produced exclusively or at higher levels in hyperproliferative cells as compared to normal cells. Target antigens include such proteins, fragments thereof and peptides which comprise at least an epitope found on such proteins. In some cases, a hyperproliferative-associated protein is the
10 product of a mutation of a gene that encodes a protein. The mutated gene encodes a protein which is nearly identical to the normal protein except it has a slightly different amino acid sequence which results in a different epitope not found on the normal protein. Such target proteins include those
15 which are proteins encoded by oncogenes such as *myb*, *myc*, *fyn*, and the translocation gene *bcr/abl*, *ras*, *src*, *P53*, *neu*, *trk* and *EGFRF*. In addition to oncogene products as target antigens, target proteins for anti-cancer treatments and protective regimens include variable regions of antibodies
20 made by B cell lymphomas and variable regions of T cell receptors of T cell lymphomas which, in some embodiments, are also used target antigens for autoimmune disease. Other tumor-associated proteins can be used as target proteins such as proteins which are found at higher levels in tumor cells
25 including the protein recognized by monoclonal antibody 17-1A and folate binding proteins.

While the present invention may be used to immunize an individual against one or more of several forms of cancer, the present invention is particularly useful to
30 prophylactically immunize an individual who is predisposed to develop a particular cancer or who has had cancer and is therefore susceptible to a relapse. Developments in genetics and technology as well as epidemiology allow for the determination of probability and risk assessment for the
35 development of cancer in individual. Using genetic screening and/or family health histories, it is possible to predict the

- 18 -

probability a particular individual has for developing any one of several types of cancer.

Similarly, those individuals who have already developed cancer and who have been treated to remove the cancer or are otherwise in remission are particularly susceptible to relapse and reoccurrence. As part of a treatment regimen, such individuals can be immunized against the cancer that they have been diagnosed as having had in order to combat a recurrence. Thus, once it is known that an individual has had a type of cancer and is at risk of a relapse, they can be immunized in order to prepare their immune system to combat any future appearance of the cancer.

The present invention provides a method of treating individuals suffering from hyperproliferative diseases. In such methods, the introduction of genetic constructs serves as an immunotherapeutic, directing and promoting the immune system of the individual to combat hyperproliferative cells that produce the target protein.

The present invention provides a method of treating individuals suffering from autoimmune diseases and disorders by conferring a broad based protective immune response against targets that are associated with autoimmunity including cell receptors and cells which produce "self"-directed antibodies.

T cell mediated autoimmune diseases include Rheumatoid arthritis (RA), multiple sclerosis (MS), Sjogren's syndrome, sarcoidosis, insulin dependent diabetes mellitus (IDDM), autoimmune thyroiditis, reactive arthritis, ankylosing spondylitis, scleroderma, polymyositis, dermatomyositis, psoriasis, vasculitis, Wegener's granulomatosis, Crohn's disease and ulcerative colitis. Each of these diseases is characterized by T cell receptors that bind to endogenous antigens and initiate the inflammatory cascade associated with autoimmune diseases. Vaccination against the variable region of the T cells would elicit an immune response including CTLs to eliminate those T cells.

In RA, several specific variable regions of T cell receptors (TCRs) which are involved in the disease have been

- 19 -

characterized. These TCRs include $V\beta$ -3, $V\beta$ -14, $V\beta$ -17 and $V\alpha$ -17. Thus, vaccination with a DNA construct that encodes at least one of these proteins will elicit an immune response that will target T cells involved in RA. See: Howell, M.D.,
5 et al., 1991 *Proc. Natl. Acad. Sci. USA* 88:10921-10925; Paliard, X., et al., 1991 *Science* 253:325-329; Williams, W.V., et al., 1992 *J. Clin. Invest.* 90:326-333; each of which is incorporated herein by reference.

In MS, several specific variable regions of TCRs
10 which are involved in the disease have been characterized. These TCRs include $V\beta$ -7 and $V\alpha$ -10. Thus, vaccination with a DNA construct that encodes at least one of these proteins will elicit an immune response that will target T cells involved in MS. See: Wucherpfennig, K.W., et al., 1990 *Science*
15 248:1016-1019; Oksenberg, J.R., et al., 1990 *Nature* 345:344-346; each of which is incorporated herein by reference.

In scleroderma, several specific variable regions of TCRs which are involved in the disease have been characterized. These TCRs include $V\beta$ -6, $V\beta$ -8, $V\beta$ -14 and $V\alpha$ -
20 16, $V\alpha$ -3C, $V\alpha$ -7, $V\alpha$ -14, $V\alpha$ -15, $V\alpha$ -16, $V\alpha$ -28 and $V\alpha$ -12. Thus, vaccination with a DNA construct that encodes at least one of these proteins will elicit an immune response that will target T cells involved in scleroderma.

In order to treat patients suffering from a T cell
25 mediated autoimmune disease, particularly those for which the variable region of the TCR has yet to be characterized, a synovial biopsy can be performed. Samples of the T cells present can be taken and the variable region of those TCRs identified using standard techniques. Genetic vaccines can
30 be prepared using this information.

B cell mediated autoimmune diseases include Lupus (SLE), Grave's disease, myasthenia gravis, autoimmune hemolytic anemia, autoimmune thrombocytopenia, asthma, cryoglobulinemia, primary biliary sclerosis and pernicious
35 anemia. Each of these diseases is characterized by antibodies which bind to endogenous antigens and initiate the inflammatory cascade associated with autoimmune diseases.

- 20 -

Vaccination against the variable region of antibodies would elicit an immune response including CTLs to eliminate those B cells that produce the antibody.

In order to treat patients suffering from a B cell mediated autoimmune disease, the variable region of the antibodies involved in the autoimmune activity must be identified. A biopsy can be performed and samples of the antibodies present at a site of inflammation can be taken. The variable region of those antibodies can be identified using standard techniques. Genetic vaccines can be prepared using this information.

In the case of SLE, one antigen is believed to be DNA. Thus, in patients to be immunized against SLE, their sera can be screened for anti-DNA antibodies and a vaccine can be prepared which includes DNA constructs that encode the variable region of such anti-DNA antibodies found in the sera.

Common structural features among the variable regions of both TCRs and antibodies are well known. The DNA sequence encoding a particular TCR or antibody can generally be found following well known methods such as those described in Kabat, et al. 1987 *Sequence of Proteins of Immunological Interest* U.S. Department of Health and Human Services, Bethesda MD, which is incorporated herein by reference. In addition, a general method for cloning functional variable regions from antibodies can be found in Chaudhary, V.K., et al., 1990 *Proc. Natl. Acad. Sci. USA* 87:1066, which is incorporated herein by reference.

In some of the embodiments of the invention that relate to gene therapy, the gene constructs contain either compensating genes or genes that encode therapeutic proteins. Examples of compensating genes include a gene which encodes dystrophin or a functional fragment, a gene to compensate for the defective gene in patients suffering from cystic fibrosis, an insulin, a gene to compensate for the defective gene in patients suffering from ADA, and a gene encoding Factor VIII. Examples of genes encoding therapeutic proteins include genes which encode erythropoietin, interferon, LDL receptor, GM-

- 21 -

CSF, IL-2, IL-4 and TNF. Additionally, genetic constructs which encode single chain antibody components which specifically bind to toxic substances can be administered.

In some preferred embodiments, the dystrophin gene is provided as part of a mini-gene and used to treat individuals suffering from muscular dystrophy. In some preferred embodiments, a mini-gene which contains coding sequence for a partial dystrophin protein is provided. Dystrophin abnormalities are responsible for both the milder Becker's Muscular Dystrophy (BMD) and the severe Duchenne's Muscular Dystrophy (DMD). In BMD dystrophin is made, but it is abnormal in either size and/or amount. The patient is mild to moderately weak. In DMD no protein is made and the patient is chair-bound by age 13 and usually dies by age 20. In some patients, particularly those suffering from BMD, partial dystrophin protein produced by expression of a mini-gene delivered according to the present invention can provide improved muscle function.

In some preferred embodiments, genes encoding IL-2, IL-4, interferon or TNF are delivered to tumor cells which are either present or removed and then reintroduced into an individual. In some embodiments, a gene encoding gamma interferon is administered to an individual suffering from multiple sclerosis.

Antisense molecules and ribozymes may also be delivered to the cells of an individual by introducing genetic material which acts as a template for copies of such active agents. These agents inactivate or otherwise interfere with the expression of genes that encode proteins whose presence is undesirable. Constructs which contain sequences that encode antisense molecules can be used to inhibit or prevent production of proteins within cells. Thus, production proteins such as oncogene products can be eliminated or reduced. Similarly, ribozymes can disrupt gene expression by selectively destroying messenger RNA before it is translated into protein. In some embodiments, cells are treated according to the invention using constructs that encode

- 22 -

antisense or ribozymes as part of a therapeutic regimen which involves administration of other therapeutics and procedures. Gene constructs encoding antisense molecules and ribozymes use similar vectors as those which are used when protein production is desired except that the coding sequence does not contain a start codon to initiate translation of RNA into protein. In some embodiments, it is preferred that the vectors described in Example 46, particularly those which contain an origin of replication and expressible form of the appropriate nuclear antigen.

Ribozymes are catalytic RNAs which are capable of self-cleavage or cleavage of another RNA molecule. Several different types of ribozymes, such as hammerhead, hairpin, Tetrahymena group I intron, axhead, and RNase P are known in the art. (S. Edgington, *Biotechnology* 1992 10, 256-262.) Hammerhead ribozymes have a catalytic site which has been mapped to a core of less than 40 nucleotides. Several ribozymes in plant viroids and satellite RNAs share a common secondary structure and certain conserved nucleotides. Although these ribozymes naturally serve as their own substrate, the enzyme domain can be targeted to another RNA substrate through base-pairing with sequences flanking the conserved cleavage site. This ability to custom design ribozymes has allowed them to be used for sequence-specific RNA cleavage (G. Paoletta et al., *EMBO* 1992, 1913-1919.) It will therefore be within the scope of one skilled in the art to use different catalytic sequences from various types of ribozymes, such as the hammerhead catalytic sequence and design them in the manner disclosed herein. Ribozymes can be designed against a variety of targets including pathogen nucleotide sequences and oncogenic sequences. Certain preferred embodiments of the invention include sufficient complementarity to specifically target the *abl-bcr* fusion transcript while maintaining efficiency of the cleavage reaction.

According to some embodiments of the present invention, cells are treated with compounds that facilitate

- 23 -

uptake of genetic constructs by the cells. According to some embodiments of the present invention, cells are treated with compounds that stimulate cell division and facilitate uptake of genetic constructs. Administration of compounds that
5 facilitate uptake of genetic constructs by the cells including cell stimulating compounds results in a more effective immune response against the target protein encoded by the genetic construct.

According to some embodiments of the present
10 invention, the genetic construct is administered to an individual using a needleless injection device. According to some embodiments of the present invention, the genetic construct is simultaneously administered to an individual intradermally, subcutaneously and intramuscularly using a
15 needleless injection device. Needleless injection devices are well known and widely available. One having ordinary skill in the art can, following the teachings herein, use needleless injection devices to deliver genetic material to cells of an individual. Needleless injection devices are well suited to
20 deliver genetic material to all tissue. They are particularly useful to deliver genetic material to skin and muscle cells. In some embodiments, a needleless injection device may be used to propel a liquid that contains DNA molecules toward the surface of the individual's skin. The liquid is propelled at
25 a sufficient velocity such that upon impact with the skin the liquid penetrates the surface of the skin, permeates the skin and muscle tissue therebeneath. Thus, the genetic material is simultaneously administered intradermally, subcutaneously and intramuscularly. In some embodiments, a needleless
30 injection device may be used to deliver genetic material to tissue of other organs in order to introduce a nucleic acid molecule to cells of that organ.

According to the invention, the genetic vaccine may be administered directly into the individual to be immunized
35 or ex vivo into removed cells of the individual which are reimplanted after administration. By either route, the genetic material is introduced into cells which are present

- 24 -

in the body of the individual. Routes of administration include, but are not limited to, intramuscular, intraperitoneal, intradermal, subcutaneous, intravenous, intraarterially, intraocularly and oral as well as
5 transdermally or by inhalation or suppository. Preferred routes of administration include intramuscular, intraperitoneal, intradermal and subcutaneous injection. Delivery of gene constructs which encode target proteins can confer mucosal immunity in individuals immunized by a mode of
10 administration in which the material is presented in tissues associated with mucosal immunity. Thus, in some examples, the gene construct is delivered by administration in the buccal cavity within the mouth of an individual.

Genetic constructs may be administered by means
15 including, but not limited to, traditional syringes, needleless injection devices, or "microprojectile bombardment gene guns". Alternatively, the genetic vaccine may be introduced by various means into cells that are removed from the individual. Such means include, for example, ex vivo
20 transfection, electroporation, microinjection and microprojectile bombardment. After the genetic construct is taken up by the cells, they are reimplanted into the individual. It is contemplated that otherwise non-immunogenic cells that have genetic constructs incorporated therein can
25 be implanted into the individual even if the vaccinated cells were originally taken from another individual.

The genetic vaccines according to the present invention comprise about 1 nanogram to about 1000 micrograms of DNA. In some preferred embodiments, the vaccines contain
30 about 10 nanograms to about 800 micrograms of DNA. In some preferred embodiments, the vaccines contain about 0.1 to about 500 micrograms of DNA. In some preferred embodiments, the vaccines contain about 1 to about 350 micrograms of DNA. In some preferred embodiments, the vaccines contain about 25 to
35 about 250 micrograms of DNA. In some preferred embodiments, the vaccines contain about 100 micrograms DNA.

- 25 -

The genetic vaccines according to the present invention are formulated according to the mode of administration to be used. One having ordinary skill in the art can readily formulate a genetic vaccine that comprises a genetic construct. In cases where intramuscular injection is the chosen mode of administration, an isotonic formulation is preferably used. Generally, additives for isotonicity can include sodium chloride, dextrose, mannitol, sorbitol and lactose. In some cases, isotonic solutions such as phosphate buffered saline are preferred. Stabilizers include gelatin and albumin. In some embodiments, a vasoconstriction agent is added to the formulation. The pharmaceutical preparations according to the present invention are provided sterile and pyrogen free.

The genetic constructs of the invention are formulated with or administered in conjunction with a polynucleotide function enhancer. Preferred co-agents according to the present invention are selected from the group consisting of benzoic acid esters, anilides, amidines, urethans and the hydrochloride salts thereof such as those of the family of local anesthetics.

The PFE may be a compound having one of the following formulae:



or



or



or



wherein:

Ar is benzene, *p*-aminobenzene, *m*-aminobenzene, *o*-aminobenzene, substituted benzene, substituted *p*-aminobenzene, substituted *m*-aminobenzene, substituted *o*-aminobenzene, wherein the amino group in the aminobenzene compounds can be amino, C₁-C₅ alkylamine, C₁-C₅, C₁-C₅ dialkylamine and

- 26 -

substitutions in substituted compounds are halogen, C₁-C₅ alkyl and C₁-C₅ alk xy;

R¹ is C=O;

R² is C₁-C₁₀ alkyl including branched alkyls;

5 R³ is hydrogen, amine, C₁-C₅ alkylamine, C₁-C₅, C₁-C₅ dialkylamine;

R² + R³ can form a cyclic alkyl, a C₁-C₁₀ alkyl substituted cyclic alkyl, a cyclic aliphatic amine, a C₁-C₁₀ alkyl substituted cyclic aliphatic amine, a heterocycle, a C₁-C₁₀ alkyl substituted heterocycle including a C₁-C₁₀ alkyl N-substituted heterocycle;

10 R⁴ is Ar, R² or C₁-C₅ alkoxy, a cyclic alkyl, a C₁-C₁₀ alkyl substituted cyclic alkyl, a cyclic aliphatic amine, a C₁-C₁₀ alkyl substituted cyclic aliphatic amine, a heterocycle, a C₁-C₁₀ alkyl substituted heterocycle and a C₁-C₁₀ alkoxy substituted heterocycle including a C₁-C₁₀ alkyl N-substituted heterocycle;

R⁵ is C=NH;

20 R⁶ is Ar, R² or C₁-C₅ alkoxy, a cyclic alkyl, a C₁-C₁₀ alkyl substituted cyclic alkyl, a cyclic aliphatic amine, a C₁-C₁₀ alkyl substituted cyclic aliphatic amine, a heterocycle, a C₁-C₁₀ alkyl substituted heterocycle and a C₁-C₁₀ alkoxy substituted heterocycle including a C₁-C₁₀ alkyl N-substituted heterocycle; and.

25 R⁷ is Ar, R² or C₁-C₅ alkoxy, a cyclic alkyl, a C₁-C₁₀ alkyl substituted cyclic alkyl, a cyclic aliphatic amine, a C₁-C₁₀ alkyl substituted cyclic aliphatic amine, a heterocycle, a C₁-C₁₀ alkyl substituted heterocycle and a C₁-C₁₀ alkoxy substituted heterocycle including a C₁-C₁₀ alkyl N-substituted heterocycle.

30 Examples of esters include: benzoic acid esters such as piperocaine, meprylcaine and isobucaine; para-aminobenzoic acid esters such as procaine, tetracaine, butethamine, propoxycaine and chloroprocaine; meta-aminobenzoic acid esters including metabuthamine and primacaine; and para-ethoxybenzoic acid esters such as parethoxycaine. Examples of anilides include lidocaine, etidocaine, mepivacaine, bupivacaine,

- 27 -

pyrrocaine and prilocaine. Other examples of such compounds include dibucaine, benzocaine, dyclonine, pramoxine, proparacaine, butacaine, benoxinate, carbocaine, methyl bupivacaine, butasin picrate, phenacaine, diothan, luccaine, 5 intracaine, nupercaine, metabutoxycaine, piridocaine, biphenamine and the botanically-derived bicyclics such as cocaine, cinnamoylcocaine, truxilline and cocaethylene and all such compounds complexed with hydrochloride.

In preferred embodiments, the PFE is bupivacaine. 10 The difference between bupivacaine and mepivacaine is that bupivacaine has a N-butyl group in place of an N-methyl group of mepivacaine. Compounds may have at that N, C₁-C₁₀. Compounds may be substituted by halogen such as procaine and chloroprocaine. The anilides are preferred.

15 Bupivacaine is administered prior to, simultaneously with or subsequent to the genetic construct. Bupivacaine and the genetic construct may be formulated in the same composition. Bupivacaine is particularly useful as a cell stimulating agent in view of its many properties and 20 activities when administered to tissue. Bupivacaine promotes and facilitates the uptake of genetic material by the cell. As such, it is a transfecting agent. Administration of genetic constructs in conjunction with bupivacaine facilitates entry of the genetic constructs into cells. Bupivacaine is 25 believed to disrupt or otherwise render the cell membrane more permeable. Cell division and replication is stimulated by bupivacaine. Accordingly, bupivacaine acts as a replicating agent. Administration of bupivacaine also irritates and damages the tissue. As such, it acts as an inflammatory agent 30 which elicits migration and chemotaxis of immune cells to the site of administration. In addition to the cells normally present at the site of administration, the cells of the immune system which migrate to the site in response to the inflammatory agent can come into contact with the administered 35 genetic material and the bupivacaine. Bupivacaine, acting as a transfection agent, is available to promote uptake of genetic material by such cells of the immune system as well.

- 28 -

Bupivacaine is related chemically and pharmacologically to the aminoacyl local anesthetics. It is a homologue of mepivacaine and related to lidocaine. Bupivacaine renders muscle tissue voltage sensitive to sodium challenge and effects ion concentration within the cells. A complete description of bupivacaine's pharmacological activities can be found in Ritchie, J.M. and N.M. Greene, *The Pharmacological Basis of Therapeutics*, Eds.: Gilman, A.G. et al, 8th Edition, Chapter 15:3111, which is incorporated herein by reference. Bupivacaine and compounds that display a functional similarity to bupivacaine are preferred in the method of the present invention.

Bupivacaine-HCl is chemically designated as 2-piperidinecarboxamide, 1-butyl-N-(2,6-dimethylphenyl)-monohydrochloride, monohydrate and is widely available commercially for pharmaceutical uses from many sources including from Astra Pharmaceutical Products Inc. (Westboro, MA) and Sanofi Winthrop Pharmaceuticals (New York, NY), Eastman Kodak (Rochester, NY). Bupivacaine is commercially formulated with and without methylparaben and with or without epinephrine. Any such formulation may be used. It is commercially available for pharmaceutical use in concentration of 0.25%, 0.5% and 0.75% which may be used on the invention. Alternative concentrations, particularly those between 0.05% - 1.0% which elicit desirable effects may be prepared if desired. According to the present invention, about 250 µg to about 10 mg of bupivacaine is administered. In some embodiments, about 250 µg to about 7.5 mg is administered. In some embodiments, about 0.05 mg to about 5.0 mg is administered. In some embodiments, about 0.5 mg to about 3.0 mg is administered. In some embodiments about 5 to 50 µg is administered. For example, in some embodiments about 50 µl to about 2 ml, preferably 50 µl to about 1500 µl and more preferably about 1 ml of 0.5% bupivacaine-HCl and 0.1% methylparaben in an isotonic pharmaceutical carrier is administered at the same site as the vaccine before, simultaneously with or after the vaccine is administered.

- 29 -

Similarly, in some embodiments, about 50 μ l to about 2 ml, preferably 50 μ l to about 1500 μ l and mor preferably about 1 ml of 0.5% bupivacaine-HCl in an isotonic pharmaceutical carrier is administered at the same site as the vaccine before, simultaneously with or after the vaccine is administered. Bupivacaine and any other similarly acting compounds, particularly those of the related family of local anesthetics may be administered at concentrations which provide the desired facilitation of uptake of genetic constructs by cells.

In some embodiments of the invention, the individual is first subject to bupivacaine injection prior to genetic vaccination by intramuscular injection. That is, up to, for example, up to a about a week to ten days prior to vaccination, the individual is first injected with bupivacaine. In some embodiments, prior to vaccination, the individual is injected with bupivacaine about 1 to 5 days before administration of the genetic construct. In some embodiments, prior to vaccination, the individual is injected with bupivacaine about 24 hrs before administration of the genetic construct. Alternatively, bupivacaine can be injected simultaneously, minutes before or after vaccination. Accordingly, bupivacaine and the genetic construct may be combined and injected simultaneously as a mixture. In some embodiments, the bupivacaine is administered after administration of the genetic construct. For example, up to about a week to ten days after administration of the genetic construct, the individual is injected with bupivacaine. In some embodiments, the individual is injected with bupivacaine about 24 hrs after vaccination. In some embodiments, the individual is injected with bupivacaine about 1 to 5 days after vaccination. In some embodiments, the individual is administered bupivacaine up to about a week to ten days after vaccination.

Other agents which may function transfecting agents and/or replicating agents and/or inflammatory agents and which may be co-administered with bupivacaine and similar acting

- 30 -

compounds include lectins, growth factors, cytokines and lymphokines such as α -interferon, gamma-interferon, platelet derived growth factor (PDGF), GC-SF, GM-CSF, TNF, epidermal growth factor (EGF), IL-1, IL-2, IL-4, IL-6, IL-8, IL-10 and 5 IL-12 as well as collagenase, fibroblast growth factor, estrogen, dexamethasone, saponins, surface active agents such as immune-stimulating complexes (ISCOMS), Freund's incomplete adjuvant, LPS analog including monophosphoryl Lipid A (MPL), muramyl peptides, quinone analogs and vesicles such as 10 squalene and squalene, hyaluronic acid and hyaluronidase may also be used administered in conjunction with the genetic construct. In some embodiments, combinations of these agents are administered in conjunction with bupivacaine and the genetic construct. For example, bupivacaine and either 15 hyaluronic acid or hyaluronidase are co-administered with a genetic construct.

The genetic construct may be combined with collagen as an emulsion and delivered parenterally. The collagen emulsion provides a means for sustained release of DNA. 50 20 μ l to 2 ml of collagen are used. About 100 μ g DNA are combined with 1 ml of collagen in a preferred embodiment using this formulation. Other sustained release formulations such as those described in *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field, which is 25 incorporated herein by reference. Such formulations include aqueous suspensions, oil solutions and suspensions, emulsions and implants as well as reservoirs and transdermal devices. In some embodiments, time release formulations for genetic constructs are preferred. In some embodiments, it is 30 preferred that the genetic construct is time released between 6-144 hours, preferably 12-96 hours, more preferably 18-72 hours.

In some embodiments of the invention, the genetic construct is injected with a needleless injection device. The 35 needleless injection devices are particularly useful for simultaneous administration of the material intramuscularly, intradermally and subcutaneously.

- 31 -

In some embodiments of the invention, the genetic construct is administered with a PFE by means of a microprojectile particle bombardment procedure as taught by Sanford et al. in U.S. Patent 4,945,050 issued July 31, 1990, which is incorporated herein by reference.

In some embodiments of the invention, the genetic construct is administered as part of a liposome complex with a polynucleotide function enhancing agent.

In some embodiments of the invention, the individual is subject to a single vaccination to produce a full, broad immune response. In some embodiments of the invention, the individual is subject to a series of vaccinations to produce a full, broad immune response. According to some embodiments of the invention, at least two and preferably four to five injections are given over a period of time. The period of time between injections may include from 24 hours apart to two weeks or longer between injections, preferably one week apart. Alternatively, at least two and up to four separate injections are given simultaneously at different sites.

In some embodiments of the invention, a complete vaccination includes injection of a single inoculant which contains a genetic construct including sequences encoding one or more targeted epitopes.

In some embodiments of the invention, a complete vaccination includes injection of two or more different inoculants into different sites. For example, in an HIV vaccine according to the invention, the vaccine comprises two inoculants in which each one comprises genetic material encoding different viral proteins. This method of vaccination allows the introduction of as much as a complete set of viral genes into the individual without the risk of assembling an infectious viral particle. Thus, an immune response against most or all of the virus can be invoked in the vaccinated individual. Injection of each inoculant is performed at different sites, preferably at a distance to ensure no cells receive both genetic constructs. As a further safety precaution, some genes may be deleted or altered to further

- 32 -

prevent the capability of infectious viral assembly. As used herein, the term "pharmaceutical kit" is meant to collectively refer to multiple inoculant used in the present invention. Such kits include separate containers containing different
5 inoculants and/or cell stimulating agents. It is intended that these kits be provided to include a set of inoculants used in an immunizing method.

The methods of the present invention are useful in the fields of both human and veterinary medicine.
10 Accordingly, the present invention relates to genetic immunization of mammals, birds and fish. The methods of the present invention can be particularly useful for mammalian species including human, bovine, ovine, porcine, equine, canine and feline species.

15 The Examples set out below include representative examples of aspects of the present invention. The Examples are not meant to limit the scope of the invention but rather serve exemplary purposes. In addition, various aspects of the invention can be summarized by the following description.
20 However, this description is not meant to limit the scope of the invention but rather to highlight various aspects of the invention. One having ordinary skill in the art can readily appreciate additional aspects and embodiments of the invention.

25 Examples

Example 1

The present invention provides an HIV vaccine using direct genetic immunization. Genetic constructs are provided which, when delivered into the cells of an individual, are
30 expressed to produce HIV proteins. According to some embodiments, the production of all viral structural proteins in the cells of the individual elicit a protective immune response which protects against HIV infection. The HIV vaccine of the present invention may be used to immunize
35 uninfected individuals from HIV infection or serve as an immunotherapeutic for those individuals already infected. The HIV vaccine of the present invention invokes an immune

- 33 -

response including CTLs which recognize and attack HIV infected cells and recognize the widest contingent of HIV protein. Thus, uninfected individuals are protected from HIV infection.

5 In some embodiments, the present invention relates to a method of immunizing an individual against HIV by administering two inoculants. These two inoculants comprise at least two and preferably more than two, a plurality or all of the genes of the HIV virus. However, the inoculants are
10 not delivered together. Accordingly, an inoculated cell will not be administered a complete complement of genes. The vaccinated individual will receive at least two different and preferably more than two, more preferably a plurality or all of the viral genes. Immune responses can then be directed at
15 the total complement of HIV protein target.

This strategy increases the probability that genetic material encoding the most effective target protein will be included in the vaccine and reduces the likelihood that a viral particle will escape detection by the immune response
20 despite structural changes in one or more viral proteins which occur when the virus undergoes mutation. Accordingly, it is desirable to vaccinate an individual with multiple and preferably a nearly complete or complete complement of genes encoding viral proteins.

25 If a single cell is provided with a complete complement of viral genes, it is possible that a complete infectious virus can be assembled within the cell. Accordingly, a genetic construct according to the present invention is not provided with such a full complement of
30 genes. Furthermore, two or more inoculants, each having an incomplete set of genes and combined having up to a full complement of viral genes, are administered to different cells, preferably at a distant site from each other to ensure that no vaccinated cell will inadvertently be exposed to a
35 full set of genes. For example, a portion of the HIV genome may be inserted into a first construct and the remaining portion of the HIV genome is inserted in a second construct.

- 34 -

The first construct is administered to an individual as a genetic vaccine in the muscle tissue of one arm while the second construct is administered to an individual as a genetic vaccine in the muscle tissue of the individual's other arm.

5 The individual may be exposed to a full set of viral genes; thus essentially vaccinating against the whole virus but with no risk that an infectious viral particle will be assembled.

As an additional safety precaution, even when genetic material is delivered by two or more inoculants at
10 distant parts of the individual's body, one or more essential genes can be deleted or intentionally altered to further ensure that an infectious viral particle cannot be formed. In such embodiments, the individual is not administered a complete functional set of viral genes.

15 A further safety precaution provides non-overlapping portions of the viral genome on the separate genetic constructs that make up the separate inoculants respectively. Accordingly, recombination between the two genetic constructs is prevented.

20 In some embodiments of the present invention, a full complement of structural genes are provided. The structural genes of HIV consist of *gag*, *pol* and *env*. These three genes are provided on two different DNA or RNA constructs. Accordingly, in one preferred embodiment, *gag* and *pol* are on
25 one DNA or RNA construct and *env* is on another. In another preferred embodiment, *gag* is on one DNA or RNA construct and *pol* and *env* is on the other. In another preferred embodiment, *gag* and *env* are on one DNA or RNA construct and *pol* is on the other. In some preferred embodiments, constructs that contain
30 *rev* have a splice acceptor upstream of the start codon for *rev*. In some preferred embodiments, constructs that contain *gag* have a splice donor upstream of the *gag* translational start codon. Optionally, in any of these combinations, HIV regulatory genes may also be present. The HIV regulatory
35 genes are: *vpr*, *vif*, *vpu*, *nef*, *tat* and *rev*.

The DNA construct in a preferred embodiment consists of a promoter, an enhancer and a polyadenylation signal. The

- 35 -

promoter may be selected from the group consisting of: HIV LTR, human Actin, human Myosin, CMV, RSV, Moloney, MMTV, human Hemoglobin, human muscle creatine and EBV. The enhancer may be selected from the group consisting of: human Actin, human Myosin, CMV, RSV, human Hemoglobin, human muscle creatine and EBV. The polyadenylation signal may be selected from the group consisting of: LTR polyadenylation signal and SV40 polyadenylation signal, particularly the SV40 minor polyadenylation signal among others.

10 In some embodiments, the two inoculant vaccine is administered intramuscularly at spatially segregated tissue of the individual, preferably in different appendages, such as for example in the right and left arms. Each inoculant of the present invention may contain from about .1 to about 1000
15 micrograms of DNA. Preferably, each inoculant contains about 1 to about 500 micrograms of DNA. More preferably, each inoculant contains about 25 to about 250 micrograms of DNA. Most preferably, each inoculant contains about 100 micrograms DNA.

20 The inoculant in some embodiments is in a sterile isotonic carrier, preferably phosphate buffered saline or saline solution.

In some embodiments, prior to vaccine administration, the tissue to be vaccinated is injected with
25 a cell proliferating agent, preferably bupivacaine. Bupivacaine injections may be performed up to about 24 hours prior to vaccination. It is contemplated that bupivacaine injection will occur immediately before vaccination. About 50 μ l to about 2 ml of 0.5% bupivacaine-HCl and 0.1%
30 methylparaben in isotonic NaCl is administered to the site where the vaccine is to be administered, preferably, 50 μ l to about 1500 μ l, more preferably about 1 ml.

In other embodiments, a cell proliferating agent, preferably bupivacaine is included in the formulation together
35 with the genetic construct. About 50 μ l to about 2 ml of 0.5% bupivacaine-HCl and 0.1% methylparaben in isotonic NaCl is administered to the site where the vaccine is to be

- 36 -

administered, preferably, 50 μ l to about 1500 μ l, more preferably about 1 ml.

Accordingly, some embodiments comprise a two inoculant vaccine: one inoculant comprising a DNA or RNA construct having two HIV structural genes, the other inoculant comprising a DNA or RNA construct having the third, remaining HIV structural gene such that the combined inoculants contain a full complement of HIV structural genes. The structural genes on each DNA construct are operably linked to a promoter, an enhancer and a polyadenylation signal. The same or different regulatory elements may control expression of the viral genes. When vaccinating an individual, the two inoculants are administered intramuscularly to different sites, preferably on different arms. In some embodiments of the invention, bupivacaine is first administered at the site where inoculant is to be administered. In some embodiments of the invention, bupivacaine is included in the formulations together with the genetic constructs.

In some embodiments, the vaccination procedure is repeated at least once and preferably two or three times. Each vaccination procedure is performed from 24 hours to two months apart.

In some embodiments, the vaccine is administered using a needleless injection device. In some embodiments, the vaccine is administered hypodermically using a needleless injection device thus providing intramuscular, intradermal, subcutaneous administration simultaneously while also administering the material interstitially.

Preferred genetic constructs include the following.

Plasmids and Cloning Strategies:

Two plasmids were constructed: one which contains HIV *gag/pol* and the other which contains HIV *env*.

The HIV-1 genomic clone pNL43 was obtained through the NIH AIDS Research and Reference Reagent Program (ARRRP), Division of AIDS, NIAID, NIH, from Dr. Malcolm Martin, and can be used as the starting material for HIV-1 viral genes for genetic constructs. Alternatively, any HIV molecular clone

- 37 -

of infected cell can, through use of the polymase chain technology, be modified sufficiently for construction including the HXB2 clone the MN clone as well as the SF or BAL-1 clone. The pNL43 clone is a construct that consists of
5 HIV-1 proviral DNA plus 3 kb of host sequence from the site of integration cloned into pUC18.

Construction of pNL-puro-env' plasmid:

This plasmid was constructed for expression of *gag* *pol*. The *StuI* site within the non-HIV 5' flanking human DNA
10 of pNL43 was destroyed by partial digestion with *StuI* followed by digestion of the free ends with *E. coli* polymerase I. The linear plasmid was filled and then self ligated, leaving a unique *StuI* site within the HIV genome. This plasmid, pNLDstu, was then digested with the blunting enzymes *StuI* and
15 *BsaBI* which eliminated a large section of the coding sequence for *gp120*. The SV40 promoter and puromycin resistance coding region (puromycin acetyl transferase (PAC)) were isolated from pBABE-puro (Morgenstern and Land, 1990 *Nucl. Acids Res.*
18(12):3587-3596, which is incorporated herein by reference,
20 kindly provided by Dr. Hartmut Land of the Imperial Cancer Research Fund) using *EcoRI* and *ClaI*. This fragment was blunted, then cloned into the *StuI/BsaBI*-digested pNLDstu. A clone was selected with the SV40-puro fragment in the correct orientation so that the 3' LTR of HIV could provide
25 poly A functions for the PAC message. This plasmid was designated pNLpuro.

Cloning strategy for deletion of *vpr* regulatory gene from the HIV *gag pol* vector:

A region from just upstream of the unique *PflMI* site
30 to just after the *vif* termination codon was amplified via PCR using primers that introduced a non-conservative amino acid change (glu->val) at amino acid 22 of *vpr*, a stop codon in the *vpr* reading frame immediately after amino acid 22, and an *EcoRI* site immediately following the new stop codon. This PCR
35 fragment was substituted for the *PflMI-EcoR I* fragment of pNLpuro or pNL43. This substitution resulted in the deletion of 122 nucleotides of the open reading frame of *vpr*, thus

- 38 -

eliminating the possibility of reversal that a point mutation strategy entails. The resulting plasmids, pNLpuroΔvpr, encode the first 21 natural amino acids of vpr plus a valine plus all other remaining HIV-1 genes and splice junctions in their native form. Such deletion strategy would also be applicable to *nef*, *vif*, and *vpu* and allow for structural gene expression but protect from the generation of a live recombinant virus. Plasmid construction for envelope expression:

The DNA segment encoding the envelope gene of HIV-1 HXB2 was cloned by the polymerase chain reaction (PCR) amplification technique utilizing the lambda cloned DNA obtained from the AIDS Research and Reference Reagent Program. The sequences of the 5' and 3' primers are 5'-AGGCGTCTCGAGACAGAGGAGAGCAAGAAATG-3' (SEQ ID NO:1) with incorporation of *Xho*I site and 5'-TTTCCTCTAGATAAGCCATCCAATCACAC-3' (SEQ ID NO: 2) with incorporation of *Xba*I site, respectively, which encompass *gp160*, *tat* and *rev* coding region. Gene specific amplification was performed using *Taq* DNA polymerase according to the manufacturer's instructions (Perkin-Elmer Cetus Corp.). The PCR reaction products were treated with 0.5 μg/ml proteinase K at 37°C for thirty minutes followed by a phenol/chloroform extraction and ethanol precipitation. Recovered DNA was then digested with *Xho*I and *Xba*I for two hours at 37°C and subjected to agarose gel electrophoresis. The isolated and purified *Xho*I-*Xba*I PCR fragment was cloned into Bluescript plasmid (Stratagene Inc., La Jolla, CA) and then subcloned into the eukaryotic expression vector pMAMneoBlue (Clontech Laboratories, Inc., Palo Alto, CA). The resulting construct was designated as pM160 (Figure 1A). The plasmid DNA was purified with CsCl gradient ultracentrifugation. The DNA construct pM160 encodes the HIV-1/HXB2 (Fisher, A.G., et al., (1985) *Nature* 316:262-265) *gp160* membrane bound glycoprotein under control of a RSV enhancer element with the MMTV LTR as a promoter.

An alternative envelope expression plasmid construction called HIV-1 env-*rev* plasmid:

- 39 -

The region encoding the two exons of rev and the vpu and envelope open reading frames of HIV-1 HXB2 was amplified via PCR and cloned into the expression vector pCND4/neo (Invitrogen). This plasmid drives envelope production through the CMV promoter.

Production and Purification:

The plasmid in *E. coli* (DH5 alpha) is grown up as follows: An LB plus ampicillin agar plate is streaked with the desired plasmid culture from frozen stock. The plate is incubated overnight (14-15 hours) at 37° C. A single colony is taken from the plate and inoculated into 15 ml of LB medium with a peptone preparation and 50 µg/ml ampicillin. This culture is grown at 37° C while being shaken (ca. 175 rpm) for 8-10 hours. OD₆₀₀ readings should be at least 1.0. 1 liter of LB medium with peptone and 50 µg/ml ampicillin is inoculated with 1.0 OD of culture. These 1-2 liter cultures are grown overnight at 37° C while being shaken (175 rpm).

Plasmid grown in *E. coli* (strain DH5 alpha) are harvested and purified by the following methods. General procedures for the lysis of cells and purification of plasmid can be found in "Molecular Cloning: A Laboratory Manual", 2nd Edition, J. Sambrook, E. F. Fritsch, and T. Maniatis, Cold Spring Harbor Press, 1989. The cells are concentrated and washed with glucose-tris-EDTA pH 8.0 buffer. The concentrated cells are lysed by treatment with lysozyme and briefly treated with 0.2 N KOH, the pH is then adjusted 5.5 with potassium acetate/acetic acid buffer. Insoluble material is removed by centrifugation. To the supernatant is added 2-propanol to precipitate the plasmid. The plasmid is redissolved in tris-EDTA buffer and further purified by phenol/chloroform extraction and an additional precipitation with 2-propanol.

Endotoxin can optionally be removed by a variety of methods including the following: specific adsorption by immobilized materials such as polymyxin (Tani et al., *Biomater. Artif. Cells Immobilization Biotechnol.* 20(2-4):457-62 (1992); Issekutz, J. *Immunol. Methods* 61(3):275-81 (1983)); anti-endotoxin monoclonal antibodies, such as 8A1 and HA-1A™

- 40 -

- (Cent c r, Malvern, PA; Bogard et al. *J. Immunol.* 150(10):4438-4449 (1993); Rietschel et al., *Infect. Immunity* page 3863 (1993)); positively charged depth filters (Hou et al., *J. Parenter. Sci. Technol.* 44(4):204-9 (Jul-Aug 1990));
- 5 poly(gamma-methyl L-glutamate), Hirayama et al., *Chem. Pharm. Bull. (Tokyo)* 40(8):2106-9 (1992)); histidine (Matsumae et al., *Biotechnol. Appl. Biochem.* 12(2):129-40 (1990)); hydrophobic interaction columns and membranes (Aida et al., *J. Immunol Methods* 132(2):191-5 (1990); Umeda et al., *Biomater*
- 10 *Artif Cells Artif Organs* 18(4):491-7 (1990); Hou et al., *Biochem. Biophys. Acta* 1073(1):149-54 (1991); Sawada et al., *J. Hyg. (London)* 97(1):103-14 (1986)); specific hydrophobic resins useful for removing endotoxin including hydrophobic polystyrene/divinylbenzene or divinylbenzene resins such as
- 15 Brownlee Polypore Resin (Applied Biosystems, Palo Alto, CA); XUS 40323.00 (Dow Chemical, Midland, MI); HP20, CHP20P (Mitsubishi Kasei, U.S.); Hamilton PRP-1, PRP-infinity (Hamilton, Reno, NV); Jordi Reversed-Phase DVB, Jordi Gel DVB, Polymer Labs PLgel™ (Alltech, Deerfield, IL); Vydac PLx™
- 20 (Separations Group, Hesperia, CA); other endotoxin removing materials and methods include Detoxi-Gel™ Endotoxin Removing Gel (Pierce Chemical, Rockford, IL); Application Note 206, (Pharmacia Biotech Inc, Piscataway, NJ). See also generally, Sharma, *Biotech. App. Biochem.* 8:5-22 (1986). Preferred anti-
- 25 endotoxin monoclonal antibodies bind to the conserved domains of endotoxin, preferably antibodies to lipid A, the most structurally conserved portion of the endotoxin molecule. Such anti-lipid A monoclonal antibodies include the high affinity murine IgG monoclonal antibody 8A1 and the human
- 30 anti-lipid A IgM(k) monoclonal antibody HA-1A™. HA-1A™ was derived from a human B *E. coli* J5 vaccine. HA-1A™. HA-1A™ is reported to be broadly cross-reactive with a variety of bacterial endotoxins (lipopolysaccharides).

Example 2

- 35 In experiments designed to compare the immunogenic response elicited by genetic vaccination and protein vaccination, animal models were designed using tumor cells

- 41 -

that specifically express a foreign target protein. Three immune competent mouse models have been developed which express foreign antigens. Three clonal tumor cell lines which are derived from the Balb/c mouse strain are used. The cell lines are: 1) a lymphoid cell line which does not metastasize significantly to other tissues but forms large palpable tumors which appear to kill the animal within an 8-12 week period; 2) a murine melanoma cell line with some ability to metastasize, mostly to the lung, and in which, following inoculation with 1 million cells, results in the development in the mice of large palpable tumors which similarly kill the animal within 10-12 weeks; and 3) a murine lung adenocarcinoma cell line which metastasizes to multiple tissues and kills the animal within 12 or more weeks. Subclones have been selected which can display foreign antigens in an unrecognized form. When transfected tumors are implanted into a parent mouse strain, unlike the majority of similar murine tumor lines, the animals do not make a protective immune response to the foreign antigens displayed and the tumors are accepted. These tumors then kill the animal with the same phenotype in the same time frame as the original untransfected tumor. Using these models, the immune response elicited by genetic vaccination against an antigen can be measured.

It was observed that mice vaccinated with a genetic vaccine comprising a genetic construct that resulted in production of the target protein by the cells of the mouse elicited an immune response including a strong cytotoxic that completely eliminated tumors displaying the target protein but with no effect on tumors that did not. In mice inoculated with the target protein itself, the immune response elicited thereby was much less effective. The tumors were reduced in size but, due to an absence of a cytotoxic response, they were not eliminated. As controls, untransfected tumors were used in experiments comparing the immune response of animals vaccinated with the genetic vaccine, subunit vaccine and unvaccinated animals. These experiments clearly demonstrate that the genetic vaccine produced a broader, more effective

- 42 -

immune response which was capable, by virtue of CTL's, of completely eliminating tumors. By contrast, immunization using intact target protein produced a more limited, less effective immune response.

5 **Example 3**

In another embodiment of the invention, a genetic vaccine against HIV has been designed. The viral protein gp160, which is processed into gp120 and gp41, is the target protein against which a genetic vaccine is produced. The
10 genetic vaccine contains a DNA construct that comprises a DNA sequence encoding gp160 operably linked regulatory elements. When administered to an individual, the DNA construct of the genetic vaccine is incorporated into the cells of the individual and gp160 is produced. The immune response that
15 is elicited by the protein is broad based and includes the humoral and both arms of the cellular immune response. The broad biological response provides superior protection to that achieved when the protein itself is administered.

Mice were injected intramuscularly with pM160,
20 described in Example 1, and subsequently analyzed for anti-HIV immune responses. The antisera from animals immunized in this manner produce anti-HIV envelope glycoprotein immune responses as measured by enzyme linked immunosorbent assay (ELISA) and immunoprecipitation assays. The antisera neutralizes HIV-1
25 infection and inhibits HIV-1 induced syncytium formation.

The observed neutralization and anti-syncytial activity may be the result of reactivity of the elicited antibodies to functionally important regions of the HIV-1 envelope protein, such as the V3 loop of gp120, CD4 binding
30 site and the N-terminal "immunodominant" region of gp41, among others.

In the genetic immunization procedure described herein, the quadriceps muscles of BALB/c mice were injected with 100 μ l of 0.5% bupivacaine-HCl and 0.1% methylparaben in
35 isotonic NaCl using a 27-gauge needle to stimulate muscle cell regeneration and facilitate uptake of the genetic construct. Twenty-four hours later, the same injection sites were then

- 43 -

injected with either 100 μ g of pM160 or with 100 μ g of pMAMneoBlue as a control plasmid (Fig. 1A). The mice were boosted by injecting the same amount of DNA construct three times at two week intervals in the same manner but without
5 pre-treatment with bupivacaine-HCl.

For the recombinant gp160 immunization, BALB/C mice were initially immunized with 1 μ g of glycosylated recombinant (HIV-1/III₈) gp160 (MicroGeneSys Inc.) in complete Freund's adjuvant followed by three boosters of 1 μ g of gp160 each in
10 incomplete Freund's adjuvant at two week intervals. The production of antibody against HIV-1 gp160 was determined by testing the mouse sera for their ability to immunoprecipitate gp160. Immunoprecipitation was performed using 1×10^6 cpm of 125 I labeled rgp160, mouse sera and protein-G agarose beads
15 (GIBCO, Inc.) as previously described by Osther, K., et al., (1991) *Hybridoma* 10:673-683, which is incorporated herein by reference. The specific precipitations were analyzed by 10% SDS-PAGE. Lane 1 is 1 μ l of preimmune mouse serum reacted with the 125 I-gp160. Lane 2 is 1 μ l of mouse serum immunized
20 from the pM160 immunized mice. Lane 3 is 1 μ l of 1:100 dilution of ID6 monoclonal anti-gp120 antibody (Ugen, K.E., et al., (1992) *Generation of Monoclonal Antibodies Against the Amino Region of gp120 Which Elicits Antibody Dependent Cellular Cytotoxicity*, Cold Spring Harbor Laboratory) as a
25 positive control. The arrow indicates the specifically immunoprecipitated 125 I-gp160 envelope glycoprotein.

125 I-labelled gp160 was specifically immunoprecipitated with antisera derived from the pM160-immunized animals (Fig. 2, lane 2) as well as the positive control anti-gp120
30 monoclonal antibody, ID6 (Fig. 2, lane 3). In contrast, the preimmune sera (Fig. 2, lane 1) only showed minimal activity in the same assay.

Eight of ten mice immunized with the pM160 construct were positive for reactivity against gp160 as determined by
35 ELISA and the immune responses from the animal with the highest anti-gp160 titer was analyzed in detail. Four mice immunized with the control vector all showed a similar

- 44 -

negative reactivity to gp160 in ELISA and one of these sera was used as the control for subsequent experiments.

It has been shown that HIV neutralizing antibodies are specifically targeted to several epitopes in gp120 and gp41, which include the V3 loop in gp120 (Goudsmit, J. et al., (1988) *AIDS* 2:157-164; and Putney, S.D., et al., (1989) *Development Of An HIV Subunit Vaccine*, Montreal), the CD4 binding site near the carboxy terminus of gp120 (Lasky, L.A., et al., (1987) *Cell* 50:975-985) as well as the immunodominant loop of gp41 just downstream of the N-terminal fusion region (Schrier, R.D., et al., (1988) *J. Virol.* 62:2531-2536).

To determine whether the anti-gp160 antibodies elicited in these mice are reactive to these important regions of the envelope glycoproteins, peptides for the BRU/V3 loop, peptides for the MN/V3 loop, peptides for the HXB2/gp41 N-terminus or peptides for HXB2/CD4 binding site were absorbed to microtiter plates and specific reactivities of the mouse antisera determined in ELISA assays. One $\mu\text{g/ml}$ of gp160 or 10 $\mu\text{g/ml}$ of each peptide was coated to microtiter plates in 0.1M bicarbonate buffer (pH 9.5) overnight at 4°C, blocked with 2% bovine serum albumin in PBS, and reacted with goat anti-mouse IgG conjugated with HRPO (Fisher) for one hour at 37°C and developed with TMB substrate (Sigma) for 10-30 minutes at room temperature in the dark. Results are reported in Figure 3. Antisera were as follows: (-+-) is preimmune sera, (-x-) is the pMAMneoBlue vector immunized sera, (-0-) is the pM160 immunized sera, (-Δ-) is from mice immunized with the rgp160 protein. Figure 3A shows results using a rgp160 protein coated plate. Figure 3B shows results using a BRU/V3 loop peptides (CNTRKRIRIQRGPGRAFTIGK (SEQ ID NO:11)) coated plate. Figure 3C shows results using a plate coated with MN/V loop peptides (YNKRKRIHIQRGPGRAFYT¹TKNIIC (SEQ ID NO:12)) with the QR sequence from HIV-1/III₂ in bold-faced type. Figure 3D shows the results using a HXB2/CD4 binding site peptides (CRIKQFINMWQEVGKAMYAPPISGIRC (SEQ ID NO:13)) coated plate. Figure 3E shows the results using a BRU/gp41 immunodominant

- 45 -

region peptides (RILAVEYIKDQQLLGIWGCSGKLIC (SEQ ID NO:14)) coated plate.

Figure 3 shows that antiserum from the pM160 construct immunized mouse has significantly higher reactivity to the BRU and MN/V3 loop peptides, the CD4 binding site peptide and the immunodominant gp41 peptide than the recombinant gp160 protein (rgp160) immunized serum. The antiserum from the rgp160 immunized mouse had much higher titer against the rgp160 than the pM160 immunized antiserum, but lower activity against the three specific neutralization epitopes of gp160 tested.

To determine whether the antisera generated by DNA immunization possessed antiviral activity, the ability of the antisera to neutralize HIV-1 infection was examined. Cell-free HIV-1/III₈ virus at 100 TCID₅₀ was incubated with serial dilutions of the antisera before being used to infect MT-2 target cells (Montefiori, D.C., (1988) *J. Clin. Microbio.* 26:231-235).

One hundred TCID₅₀ HIV-1/III₈ cell-free virus was preincubated with serial dilutions of antisera for one hour at 37°C. Following incubation the pretreated virus was then plated on the 4x10⁴ of target cell line, MT-2 for one hour at 37°C, following infection the MT-2 cells were washed three times and then incubated at 37°C at 5% CO₂. Fusion was evaluated three days later quantitatively by visually counting the number of syncytia per well in triplicate experiments under a phase contrast microscope.

The results are reported in Figure 4. Figure 4A shows the results using vector-immunized mouse sera compared with Figure 4B which shows the results using pM160 immunized sera. Neutralization values (V_n/V_o) versus the dilution factors (Nara, P., (1989) *Techniques In HIV Research* eds. Aldovini, A. & Walkter, B.D., 77-86 M Stockton Press) are illustrated in Figure 4C. The control serum (-x) was from pMAMneoBlue vector immunized mice. The test sera (O) were from pM160 immunized mice.

- 46 -

Syncytia inhibition was performed as described by Osth r, K., et al., (1991) *Hybridoma* 10:673-683. The H9/III₈ c ll line was pre-incubated with serial dilutions (1:100, 1:200, and 1:400) of antisera were made in 96 well plates in a total volume of 50 μ l for thirty minutes at 37°C at 5% CO₂. Fusion was evaluated three days later quantitatively by visually counting the number of syncytia per well under a phase construct microscope. Figure 4D is the target cells co-cultivated with HIV-1/III₈ cell line treated with preimmune serum. Figure 4E is the same as Figure 4D but treated with vector control immunized serum. Figure 4F is the same as Figure 4D but treated with rgp160 immunized serum. Figure 4G is the same as Figure 4D but treated with pM160 immunized serum. Figures 4D to 4G show that inhibition of syncytia was apparent at dilution at 1:200 in these assays. MT-2 cells were infected with cell-free HIV-1/III₈ which had been preincubated with vector-immunized antiserum readily formed syncytia (Figure 4A). In comparison, preincubation with pM160 immunized mouse serum prevented syncytium formation (Figure 4B). The neutralization kinetics were determined by V_n/V_0 versus serial dilutions of antisera (Nara, P., (1989) *Techniques In HIV Research*, eds. Aldovini, A. & Walker, B.D., 77-86, M Stockton Press) (Figure 4C). The serum from the pM160 immunized mouse had biologically active neutralizing activity at dilutions of up to 1:320 while control antisera did not show similar activity.

To determine if the antiserum from the pM160 immunized mouse could inhibit envelope-mediated virus spread through direct cell-to-cell fusion, syncytium inhibition assays were performed. Antiserum from the pM160 immunized mouse inhibits HIV-1 induced syncytium formation at 1:200 dilutions (Fig. 4G). In contrast, the preimmune sera (Fig. 4D), antisera from the rgp160 immunized mice (Fig. 4F) and antisera from the control vector-immunized animals (Fig. 4E) failed to inhibit syncytium formation at the same dilutions.

Observations from the neutralization (Figures 4A-C) and syncytium inhibition assays (Figures D-G) of these sera

- 47 -

correlates with the observed ELISA reactivities (Fig. 3). The antiserum from the pM160 immunized mouse which showed a high level of binding to neutralizing epitopes likewise demonstrated high level anti-viral activities; conversely, sera with little binding to these epitopes including the antiserum from rgp160 immunized mice have low anti-viral activity.

To test whether the antisera from pM160 immunized mice can inhibit gp120 binding to CD4-bearing T-cells, a direct inhibition assay monitored by fluorocytometry was employed (Chen, Y.H., et al., (1992) *AIDS* 6:533-539. It was observed that serum from the pM160 construct-immunized mouse was able to block the binding of gp120 to CD4-bearing T-cells: a 1:15 dilution of immune serum inhibited FITC-gp120 binding to CD4⁺SupT1 cells by 22% \pm 2% in replicate experiments as evaluated by flow cytometry. This indicates that this region for HIV entry into target cells can also be functionally inhibited by this antiserum. These data are consistent with observed ELISA reactivity of the antiserum to the CD4 binding site peptides (Fig. 3c).

Immunoglobulin isotyping studies were performed by using a commercial murine monoclonal antibody isotyping kit (Sigma). Of the anti-gp160 specific antibodies elicited by pM160 immunization, 19% are IgG1, 51% are IgG2, 16% are IgG3, 10% are IgM and 5% are IgA. The predominance of IgG isotypes indicates that a secondary immune response has taken place, and further suggests that helper T-cells can be elicited by genetic immunization.

pM160 and pMAMneoBlue DNAs were coated onto microtiter plates and specific binding was determined by ELISA using sera all immunized animals. No significant binding to plasmid DNA was observed. Thus, using genetic material for inoculation into muscle tissue appears unlikely to produce an anti-plasmid DNA response.

Introducing construct DNA into mouse muscle by needle injection may cause inconsistent results, as this technique does not provide a means to control DNA uptake by

- 48 -

muscle cells. Injection of construct DNA alone (n≈4) with bupivacaine pretreated animals (n≈4) was compared. The immun responses observed in the two groups were dissimilar, with 25% and 75% animals responding in ELISA assays respectively.

- 5 Increased efficiency may be achieved by use of a direct DNA delivery system such as particle bombardment (Klein, T.M. et al., (1992) *Bio/technology* 10:286-291.

Evidence of neutralization, syncytia inhibition, inhibition of CD4-gp120 binding, and specific binding to
10 several important regions on the gp160 demonstrate that introduction of a DNA construct encoding HIV gp160 membrane-bound glycoprotein directly into muscle cells of living animals can elicit specific humoral responses, and generate biologically relevant anti-viral antibodies.

- 15 To test whether the vaccine is capable of eliciting a protective immune response, the animal model described above was used. Tumor cells were transfected with DNA encoding p160, confirmed to express the protein and implanted into the animal. Controls included untransfected tumor lines.

- 20 Genetically immunized animals were vaccinated with plasmid pm160. Controls included unvaccinated animals, animals vaccinated with vector DNA only and animals administered the gp160 protein.

- Results demonstrate that the immune response of
25 genetically vaccinated mice was sufficient to completely eliminate the transfected tumors while having no effect on untransfected tumors. gp160 protein vaccination led to some reduction in tumor size in transfected tumors as compared to untransfected tumors but had no effect on mortality.
30 Unvaccinated animals showed similar mortality for both transfected and untransfected tumors.

Example 4

- The following is a list of constructs which may be used in the methods of the present invention. The vector
35 pBabe.puro, which is used as a starting material to produce many of the below listed constructs, was originally constructed and reported by Morgenstern, J.P. and H. Land,

- 49 -

1990 *Nucl. Acids Res.* 18(12):3587-3596, which is incorporated herein by reference. The pBabe.puro plasmid is particularly useful for expression of exogenous genes in mammalian cells. DNA sequences to be expressed are inserted at cloning sites
5 under the control of the Moloney murine leukemia virus (Mo MuLV) long terminal repeat (LTR) promoter. The plasmid contains the selectable marker for puromycin resistance.

Example 5

Plasmid pBa.Va3 is a 7.8 kb plasmid that contains
10 a 2.7 kb *EcoRI* genomic fragment encoding the T cell receptor Va3 region containing the L, V and J segments cloned into the *EcoRI* site of pBabe.puro. The T cell receptor-derived target protein is useful in the immunization against and treatment of T cell mediated autoimmune disease and clonotypic T cell
15 lymphoma and leukemia.

Example 6

Plasmid pBa.gagpol-vpr is a 9.88 kb plasmid that contains the *gag/pol* and *vif* genes from HIV/MN cloned into pBabe.puro. The *vpr* gene is deleted. The plasmid which
20 contains these HIV viral genes, which encode HIV target proteins, is useful in the immunization against and treatment of HIV infection and AIDS. The HIV DNA sequence is published in Reiz, M.S., 1992 *AIDS Res. Human Retro.* 8:1549, which is incorporated herein by reference. The sequence is accessible
25 from Genbank No.: M17449, which is incorporated herein by reference.

Example 7

Plasmid pM160 is an 11.0 kb plasmid that contains the 2.3 kb PCR fragment encoding the HIV-I/3B envelope protein
30 and *rev/tat* genes cloned into pMAMneoBlue. The *nef* region is deleted. The plasmid which contains these HIV viral genes, which encode HIV target proteins, is useful in the immunization against and treatment of HIV infection and AIDS. The DNA sequence of HIV-1/3B is published in Fisher, A., 1985
35 *Nature* 316:262, which is incorporated herein by reference. The sequence is accessible from Genbank No.: K03455, which is incorporated herein by reference.

- 50 -

Example 8

Plasmid pBa.VL is a 5.4 kb plasmid that contains PCR fragment encoding the VL region of an anti-DNA antibody cloned into pBabe.puro at the *Xba*I and *Eco*RI sites. The antibody-derived target protein is an example of a target protein useful in the immunization against and treatment of B cell mediated autoimmune disease and clonotypic B cell lymphoma and leukemia. A general method for cloning functional variable regions from antibodies can be found in Chaudhary, V.K., et al., 1990 *Proc. Natl. Acad. Sci. USA* 87:1066, which is incorporated herein by reference.

Example 9

Plasmid pOspA.B is a 6.84 kb plasmid which contains the coding regions encoding the OspA and OspB antigens of the *Borrelia burgdorferi*, the spirochete responsible for Lyme's disease cloned into pBabe.puro at the *Bam*HI and *Sal*I sites. The PCR primers used to generate the OspA and OspB fragments are 5'-GAAGGATCCATGAAAAATATTTATTGGG-3' (SEQ ID NO:3) and 5'-ACTGTCGACTTATTTTAAAGCGTTTTTAAG-3' (SEQ ID NO: 4). See: Williams, W.V., et al. 1992 *DNA and Cell. Biol.* 11(3):207, which is incorporated herein by reference. The plasmid which contains these pathogen genes, which encode target proteins, is useful in the immunization against Lyme's disease.

Example 10

Plasmid pBa.Rb-G is a 7.10 kb plasmid which contains a PCR generated fragment encoding the rabies G protein cloned into pBabe.puro at the *Bam*HI site. The plasmid which contains this pathogen gene, which encodes the rabies G protein, is useful in the immunization against Rabies. The DNA sequence is disclosed in Genebank No.:M32751, which is incorporated herein by reference. See also: Anilionis, A., et al., 1981 *Nature* 294:275, which is incorporated herein by reference.

Example 11

Plasmid pBa.HPV-L1 is a 6.80 kb plasmid which contains a PCR generated fragment encoding the L1 capsid protein of the human papillomavirus (HPV) including HPV strains 16, 18, 31 and 33 cloned into pBabe.puro at the *Bam*HI

- 51 -

and *EcoRI* sites. The plasmid is useful in the immunization against HPV infection and the cancer caused thereby. The DNA sequence is disclosed in Genbank No.:M15781, which is incorporated herein by reference. See also: Howley, P., 1990 *Fields Virology*, Volume 2, Eds.: Channock, R.M. et al. Chapter 58:1625; and Shah, K. and P. Howley, 1990 *Fields Virology*, Volume 2, Eds.: Channock, R.M. et al. Chapter 59; both of which are incorporated herein by reference.

Example 12

10 Plasmid pBa.HPV-L2 is a 6.80 kb plasmid which contains a PCR generated fragment encoding the L2 capsid protein of the human papillomavirus (HPV) including HPV strains 16, 18, 31 and 33 cloned into pBabe.puro at the *BamHI* and *EcoRI* sites. The plasmid is useful in the immunization
15 against HPV infection and the cancer caused thereby. The DNA sequence is disclosed in Genbank No.:M15781, which is incorporated herein by reference. See also: Howley, P., 1990 *Fields Virology*, Volume 2, Eds.: Channock, R.M. et al. Chapter 58:1625; and Shah, K. and P. Howley, 1990 *Fields Virology*,
20 Volume 2, Eds.: Channock, R.M. et al. Chapter 59; both of which are incorporated herein by reference.

Example 13

Plasmid pBa.MNp7 is a 5.24 kb plasmid which contains a PCR generated fragment encoding the p7 coding region
25 including the HIV MN gag (core protein) sequence cloned into pBabe.puro at the *BamHI* site. The plasmid which contains these HIV viral genes, which encode HIV target proteins, is useful in the immunization against and treatment of HIV infection and AIDS. Reiz, M.S., 1992 *AIDS Res. Human Retro.*
30 8:1549, which is incorporated herein by reference. The sequence is accessible from Genbank No.:M17449, which is incorporated herein by reference.

Example 14

Plasmid pGA733-2 is a 6.3 kb plasmid that contains
35 the GA733-2 tumor surface antigen cloned from the colorectal carcinoma cell line SW948 into pCDM8 vector (Seed, B. and A. Aruffo, 1987 *Proc. Natl. Acad. Sci. USA* 84:3365, which is

- 52 -

incorporated herein by reference) at *Bst*XI site. The tumor-associated target protein is an example of a target protein useful in the immunization against and treatment of hyperproliferative disease such as cancer. The GA733-2 antigen is a useful target antigen against colon cancer. The GA733 antigen is reported in Szala, S. et al., 1990 *Proc. Natl. Acad. Sci. USA* 87:3542-3546, which is incorporated herein by reference.

Example 15

Plasmid pT4-pMV7 is a 11.15 kb plasmid that contains cDNA which encodes human CD4 receptor cloned into pMV7 vector at the *Eco*RI site. The CD4 target protein is useful in the immunization against and treatment of T cell lymphoma. Plasmid pT4-pMV7 is available from the AIDS Repository, Catalog No. 158.

Example 16

Plasmid pDJGA733 is a 5.1 kb plasmid that contains the GA733 tumor surface antigen cloned into pBabe.puro at the *Bam*HI site. The tumor-associated target protein is an example of a target protein useful in the immunization against and treatment of hyperproliferative disease such as cancer. The GA733 antigen is a useful target antigen against colon cancer.

Example 17

Plasmid pBa.RAS is a 6.8 kb plasmid that contains the ras coding region that was first subcloned from pZIPneoRAS and cloned into pBabe.puro at the *Bam*HI site. The ras target protein is an example of a cytoplasmic signalling molecule. The method of cloning ras is reported in Weinberg 1984 *Mol. Cell. Biol.* 4:1577, which is incorporated herein by reference. Ras encoding plasmid are useful for the immunization against and treatment of hyperproliferative disease such as cancer; in particular, ras related cancer such as bladder, muscle, lung, brain and bone cancer.

Example 18

Plasmid pBa.MNp55 is a 6.38 kb plasmid which contains a PCR generated fragment encoding the p55 coding region including the HIV MN gag precursor (core protein)

- 53 -

sequence cloned into pBabe.puro at the *Bam*HI site. The plasmid which contains these HIV viral genes, which encode HIV target proteins, is useful in the immunization against and treatment of HIV infection and AIDS. Reiz, M.S., 1992 *AIDS Res. Human Retro.* 8:1549, which is incorporated herein by reference. The sequence is accessible from Genbank No.: M17449, which is incorporated herein by reference.

Example 19

Plasmid pBa.MNp24 is a 5.78 kb plasmid which contains a PCR generated fragment from the pMN-SF1 template encoding the p24 coding region including the whole HIV MN gag coding region cloned into pBabe.puro at the *Bam*HI and *Eco*RI sites. The plasmid which contains these HIV viral genes, which encode HIV target proteins, is useful in the immunization against and treatment of HIV infection and AIDS. Reiz, M.S., 1992 *AIDS Res. Human Retro.* 8:1549, which is incorporated herein by reference. The sequence is accessible from Genbank No.: M17449, which is incorporated herein by reference.

Example 20

Plasmid pBa.MNp17 is a 5.5 kb plasmid which contains a PCR generated fragment encoding the p17 coding region including the HIV MN gag (core protein) sequence cloned into pBabe.puro at the *Bam*HI and *Eco*RI sites. The plasmid which contains these HIV viral genes, which encode HIV target proteins, is useful in the immunization against and treatment of HIV infection and AIDS. Reiz, M.S., 1992 *AIDS Res. Human Retro.* 8:1549, which is incorporated herein by reference. The sequence is accessible from Genbank No.: M17449, which is incorporated herein by reference.

Example 21

Plasmid pBa.SIVenv is a 7.8 kb plasmid which contains a 2.71 PCR generated fragment amplified from a construct containing SIV 239 in pBR322 cloned into pBabe.puro at the *Bam*HI and *Eco*RI sites. The primers used are 5'-GCCAGTTTTGGATCCTTAAAAAAGGCTTGG-3' (SEQ ID NO:5) and 5'-TTGTGAGGGACAGAATTCCAATCAGGG-3' (SEQ ID NO:6). The plasmid is

- 54 -

available from the AIDS Research and Reference Reagent Program; Catalog No. 210.

Example 22

Plasmid pcTSP/ATK.env is a 8.92 kb plasmid which
5 contains a PCR generated fragment encoding the complete HTLV
envelope coding region from HTLV-1/TSP and /ATK isolates
subcloned into the pcDNA1/neo vector. The primers used are 5'-
CAGTGATATCCCGGGAGACTCCTC-3' (SEQ ID NO:7) and 5'-
GAATAGAAGAACTCCTCTAGAATTC-3' (SEQ ID NO:8). Plasmid
10 pcTSP/ATK.env is reported in 1988 *Proc. Natl. Acad. Sci. USA*
85:3599, which is incorporated herein by reference. The HTLV
env target protein is useful in the immunization against and
treatment of infection by HTLV and T cell lymphoma.

Example 23

15 Plasmid pBa.MNgp160 is a 7.9 kb plasmid which
contains a 2.8 kb PCR generated fragment amplified from a
construct containing MNenv in pSP72 and cloned into pBabe.puro
at the *Bam*HI and *Eco*RI sites. The primers used are 5'-
GCCTTAGGCGGATCCTATGGCAGGAAG-3' (SEQ ID NO:9) and 5'-
20 TAAGATGGGTGGCCATGGTGAATT-3' (SEQ ID NO:10). Reiz, M.S., 1992
AIDS Res. Human Retro. 8:1549, which is incorporated herein
by reference. The sequence is accessible from Genbank
No.:M17449, which is incorporated herein by reference. The
plasmid which contains these HIV viral genes, which encode HIV
25 target proteins, is useful in the immunization against and
treatment of HIV infection and AIDS.

Example 24

Plasmid pC.MNp55 is a 11.8 kb plasmid which contains
a 1.4 kb PCR generated fragment amplified from the *gag* region
30 of MN isolate and cloned into the pCEP4 vector. The plasmid
which contains these HIV viral genes, which encode HIV target
proteins, is useful in the immunization against and treatment
of HIV infection and AIDS. Reiz, M.S., 1992 *AIDS Res. Human*
Retro. 8:1549, which is incorporated herein by reference. The
35 sequence is accessible from Genbank No.: M17449, which is
incorporated herein by reference.

Example 25

- 55 -

Plasmid pC.Neu is a 14.2 kb plasmid that contains a 3.8 kb DNA fragment containing the human neu oncogene coding region that was cut out from the LTR-2/erbB-2 construct and subcloned into the pCEP4 vector. The pC.Neu plasmid is reported in DiFiore 1987 *Science* 237:178, which is incorporated herein by reference. The neu oncogene target protein is an example of a growth factor receptor useful as a target protein for the immunization against and treatment of hyperproliferative disease such as cancer; in particular, colon, breast, lung and brain cancer.

Example 26

Plasmid pC.RAS is a 11.7 kb plasmid that contains a 1.4 kb DNA fragment containing the ras oncogene coding region that was first subcloned from pZIPneoRAS and subcloned into pCEP4 at the BamHI site. The pC.RAS plasmid is reported in Weinberg 1984 *Mol. Cell. Biol.* 4:1577, which is incorporated herein by reference. The ras target protein is an example of a cytoplasmic signalling molecule. Ras encoding plasmid are useful for the immunization against and treatment of hyperproliferative disease such as cancer; in particular, ras related cancer such as bladder, muscle, lung, brain and bone cancer.

Example 27

Plasmid pNLpuro is a 15 kb plasmid which contains HIV gag/pol and SV40-puro insertion. The plasmid which contains these HIV viral genes which encode HIV target proteins, is useful in the immunization against and treatment of HIV infection and AIDS.

Example 28

A DNA construct was designed to test the effectiveness of a genetic vaccine against human CD4 in mice. These experiments were designed to test the ability of a vaccine to protect against a T lymphoma antigen. In T cell lymphoma, CD4 is a tumor specific antigen. Accordingly, this model demonstrates the ability of the genetic vaccine to protect against T lymphoma. Further, these experiments tested the effectiveness against a member of the immunoglobulin

- 56 -

superfamily of molecules. CD4 is highly conserved between human and murine species.

The animal model used was described above. Tumor cells were transfected with DNA encoding CD4, confirmed to
5 express the protein and implanted into the animal. Controls included untransfected tumor lines. Although the animals were immunocompetent, an immune response was not directed against the implanted, CD4-labelled tumors in unvaccinated animals.

Genetically immunized animals were vaccinated with
10 plasmid pT4-pMV7, described in Example 15. Controls included unvaccinated animals and animals administered the CD4 protein.

In the genetic immunization procedure described herein, the quadriceps muscles of BALB/c mice were injected with 100 μ l of 0.5% bupivacaine-HCl and 0.1% methylparaben in
15 isotonic NaCl using a 27-gauge needle to stimulate muscle cell regeneration to facilitate uptake of the genetic construct. Twenty-four hours later, the same injection sites were then injected with either 100 μ g of pT4-pMV7 or with 100 μ g of pMV7 as a control plasmid. The mice were boosted by injecting the
20 same amount of DNA construct three times at two week intervals in the same manner but without pre-treatment with bupivacaine-HCl.

Animals received 1,000,000 CD4-labelled tumor cells. In non-vaccinated animals, large tumors formed and death
25 resulted after about 7-10 weeks. Vaccinated animals did not develop similar deadly tumors.

Results demonstrate that the immune response of genetically vaccinated mice was sufficient to completely eliminate the transfected tumors while having no effect on
30 untransfected tumors. CD4 protein vaccination led to some reduction in tumor size in transfected tumors as compared to untransfected tumors but had no effect on mortality. Unvaccinated animals showed similar mortality for both transfected and untransfected tumors.

35 Example 29

A DNA construct was designed to test the effectiveness of a genetic vaccine against human GA733 in

- 57 -

mice. These experiments were designed to test the ability of a vaccine to protect against GA733 associated cancer such as colon cancer. The animal model used was described above. Tumor cells were transfected with DNA encoding GA733, confirmed to express the protein and implanted into the animal. Controls included untransfected tumor lines.

Genetically immunized animals were vaccinated with plasmid pGA733-2, described in Example 14, following the method described above. Controls included unvaccinated animals and animals administered the GA733 protein.

Results demonstrate that the immune response of genetically vaccinated mice was sufficient to completely eliminate the transfected tumors while having no effect on untransfected tumors. GA733 protein vaccination led to some reduction in tumor size in transfected tumors as compared to untransfected tumors but had no effect on mortality. Unvaccinated animals showed similar mortality for both transfected and untransfected tumors.

Example 30

A DNA construct was designed to test the effectiveness of a genetic vaccine against human p185neu in mice. These experiments were designed to test the ability of a vaccine to protect against p185neu associated cancer such as breast, lung and brain cancer. The animal model used was described above. Tumor cells were transfected with DNA encoding neu, confirmed to express the protein and implanted into the animal. Controls included untransfected tumor lines.

Genetically immunized animals were vaccinated with plasmid pLTR-2/erbB-2, a 14.3 kb plasmid that contains the human neu oncogene coding region cloned into the LTR-2 vector at the XhoI site following the method described above. The 5'LTR and 3'LTR are from Moloney-MuLV LTR. Controls included unvaccinated animals and animals administered the p185neu protein.

Results demonstrate that the immune response of genetically vaccinated mice was sufficient to completely eliminate the transfected tumors while having no effect on

- 58 -

untranslated tumors. p185 protein vaccination led to some reduction in tumor size in transfected tumors as compared to untransfected tumors but had no effect on mortality. Unvaccinated animals showed similar mortality for both transfected and untransfected tumors.

Example 31

A DNA construct was designed to test the effectiveness of a genetic vaccine against human Ras in mice. These experiments were designed to test the ability of a vaccine to protect against Ras associated cancer such as bladder, muscle, lung, brain and bone cancer. The animal model used was described above. Tumor cells were transfected with DNA encoding Ras, confirmed to express the protein and implanted into the animal. Controls included untransfected tumor lines.

Genetically immunized animals were vaccinated with plasmid pBa.RAS, described in Example 17 following the vaccination method described above. The ras target protein is an example of a cytoplasmic signalling molecule. The method of cloning ras is reported in Weinberg 1984 *Mol. Cell. Biol.* 4:1577, which is incorporated herein by reference. Controls included unvaccinated animals and animals administered the Ras protein.

Example 32

A DNA construct was designed to test the effectiveness of a genetic vaccine against human rabies G protein antigen in mice. The animal model used was described above. Tumor cells were transfected with DNA encoding rabies G protein, confirmed to express the protein and implanted into the animal. Controls included untransfected tumor lines.

Genetically immunized animals were vaccinated with plasmid pBa.Rb-G which is described in Example 10, following the vaccination method described above. The rabies G target protein is an example of a pathogen antigen. The DNA sequence is disclosed in Genbank No.:M32751. Controls included unvaccinated animals and animals administered the G protein.

Example 33

- 59 -

A DNA construct was designed to test the effectiveness of a genetic vaccine against Lyme's disease antigen in mice. The animal model used was described above. Tumor cells were transfected with DNA encoding OspA and Osp
5 B, confirmed to express the protein and implanted into the animal. Controls included untransfected tumor lines.

Genetically immunized animals were vaccinated with plasmid pOspA.B which is described in Example 9. Controls included unvaccinated animals and animals administered OspA
10 and OspB proteins.

Example 34

A DNA construct was designed to test the effectiveness of a genetic vaccine against a human T cell receptor variable region in mice. These experiments were
15 designed to test the ability of a vaccine to protect against a T cell receptor derived protein associated cancer such as T cell lymphoma and T cell mediated autoimmune disease. The animal model used was described above. Tumor cells were transfected with DNA encoding Ras, confirmed to express the
20 protein and implanted into the animal. Controls included untransfected tumor lines.

Genetically immunized animals were vaccinated with plasmid pBa.Va3 which is described in Example 5 following the vaccination method described above.

25 Example 35

The plasmid pM160 can be used as a starting material for several plasmids useful to express one or more genes from the env portion of HIV. Construction of pM160 is described above. The plasmid encompasses gp160, tat and rev coding
30 region. The nef gene is absent.

The promoter controlling gp160/rev/tat gene expression is MMTV LTR. The promoter may be deleted and replaced with Actin promoter, myosin promoter, HIV LTR promoter and CMV promoter.

35 The gene conferring ampicillin resistance may be deleted or otherwise inactivated. The gene conferring

- 60 -

neomycin resistance may be placed under the control of a bacterial promoter.

The Rous sarcoma virus enhancer may be deleted from the plasmid. The RSV enhancer may be replaced with the muscle
5 creatine enhancer.

The gp160/rev/tat genes overlap and share the same nucleotide sequences in different reading frames. The rev gene may be deleted by changing its initiation codon to a different codon. Similarly, the tat gene may be eliminated
10 by the same means. In each plasmid except those using the HIV LTR promoter to control gp160/rev/tat, either rev, tat, or both rev and tat may be eliminated. In plasmids using the HIV LTR promoter, tat must be present.

The following Table lists pM160-modified plasmids.
15 Each plasmid has an inactivated ampicillin gene. Each has deleted the RSV enhancer. Some have no enhancer (no); some have creatine muscle enhancer (CME). Some have the HIV rev gene (yes) while it is deleted in others (no). Some have the HIV tat gene (yes) while it is deleted in others (no).

Construct	Promoter	enhancer	rev	tat
20 RA-1	Actin	no	yes	yes
RA-2	Actin	no	yes	no
RA-3	Actin	no	no	yes
RA-4	Actin	CME	yes	yes
25 RA-5	Actin	CME	yes	no
RA-6	Actin	CME	no	yes
RA-7	CMV	no	yes	yes
RA-8	CMV	no	yes	no
RA-9	CMV	no	no	yes
30 RA-10	CMV	CME	yes	yes
RA-11	CMV	CME	yes	no
RA-12	CMV	CME	no	yes
RA-13	MMTV	no	yes	yes
RA-14	MMTV	no	yes	no
35 RA-15	MMTV	no	no	yes
RA-16	MMTV	CME	yes	yes
RA-17	MMTV	CME	yes	no
RA-18	MMTV	CME	no	yes
RA-19	Myosin	no	yes	yes
40 RA-20	Myosin	no	yes	no
RA-21	Myosin	no	no	yes
RA-22	Myosin	CME	yes	yes
RA-23	Myosin	CME	yes	no
RA-24	Myosin	CME	no	yes
45 RA-25	HIV-1 LTR	no	yes	yes
RA-26	HIV-1 LTR	no	no	yes

- 61 -

RA-27	HIV-1 LTR CME	yes	yes
RA-28	HIV-1 LTR CME	no	yes

Constructions RA-29 to RA-56 are identical to RA-1 to RA-32 respectively except in each case the promoter
5 controlling the neomycin gene is a bacterial promoter.

Example 36

The plasmid pNLpuro may be used as a starting material to produce several different plasmids which express the HIV *gag/pol* genes. As described above, pNLpuro was
10 constructed for expression of *gag pol*. The plasmid pNLpuroΔvpr, which is described above, was designed to delete the vpr regulatory gene from the HIV *gag pol* vector in order to eliminate a necessary regulatory protein from the set of genes to be introduced by vaccination. In addition to vpr,
15 other changes may be made by those having ordinary skill in the art to plasmid pNL43puro using standard molecular biology techniques and widely available starting material.

The human flanking sequences 5' and 3' of the HIV sequences can be removed by several methods. For example,
20 using PCR, only HIV, SV40-puro, and pUC18 sequences can be amplified and reconstructed.

The *psi* region of HIV, which is important in the packaging of the virus, can be deleted from pNL43puro-based plasmids. In order to delete the *psi* region, the pNLpuro
25 plasmid is cut with *SacI* and *SpeI*. This digestion removes the *psi* region as well as the 5' LTR which is upstream and portion of the *gag/pol* region which is downstream of *psi*. In order to reinsert the deleted non-*psi* sequences, PCR amplification is performed to regenerate those sequences.
30 Primers are designed which regenerate the portions of the HIV sequence 5' and 3' to *psi* without regenerating *psi*. The primers reform the *SacI* site at the portion of the plasmid 5' of the 5' LTR. Primers go downstream from a site upstream of the *SacI* site to a site just 3' of the 5' end of the *psi*
35 region, generating an *AatI* site at the 3' end. Primers starting just 5' of the *psi* region also generate an *AatI* site and, starting 3' of the *SpeI* site, regenerate that site. The

- 62 -

PCR generated fragments are digested with *SacI*, *AatI* and *SpeI* and ligated together with the *SacI/SpeI* digested pNLpuro-*psi* fragment. The HIV 5'LTR promoter can be deleted and replaced with Moloney virus promoter, MMTV LTR, Actin promoter, myosin promoter and CMV promoter.

The HIV 3'LTR polyadenylation site can be deleted and replaced with SV40 polyadenylation site.

The gene conferring ampicillin resistance may be deleted or otherwise inactivated.

The following is a list of pNLpuro-based constructions in which HIV *psi* and *vpr* regions are deleted and human flanking regions 5' and 3' of the HIV sequences are deleted.

	Construct	Promoter	poly(A)	Amp ^r
15	LA-1	Moloney	HIV 3'LTR	yes
	LA-2	Moloney	SV40	yes
	LA-3	Moloney	HIV 3'LTR	no
	LA-4	Moloney	SV40	no
	LA-5	CMV	HIV 3'LTR	yes
20	LA-6	CMV	SV40	yes
	LA-7	CMV	HIV 3'LTR	no
	LA-8	CMV	SV40	no
	LA-9	MMTV	HIV 3'LTR	yes
	LA-10	MMTV	SV40	yes
25	LA-11	MMTV	HIV 3'LTR	no
	LA-12	MMTV	SV40	no
	LA-13	HIV 5' LTR	HIV 3'LTR	yes
	LA-14	HIV 5' LTR	SV40	yes
	LA-15	HIV 5' LTR	HIV 3'LTR	no
30	LA-16	HIV 5' LTR	SV40	no

Constructions LA-17 to LA-32 are identical to LA-1 to LA-16 respectively except in each case at least one of the human flanking sequence remains.

Example 37

In another construction for expressing the *env* gene, that region of HIV may be inserted into the commercially available plasmid pCEP4 (Invitrogen). The pCEP4 plasmid is particularly useful since it contains the Epstein Barr virus origin of replication and nuclear antigen EBNA-1 coding region which produces high copy episomal replication without integration. pCEP4 also contains the hygromycin marker under the regulatory control of the thymidine kinase promoter and

- 63 -

polyadenylation site. The HIV *env* coding region is placed under the regulatory control of the CMV promoter and SV40 polyadenylation site. The HIV *env* coding region was obtained as a 2.3 kb PCR fragment from HIV/3B, Genbank sequence
5 K03455. The resulting pCEP4-based plasmid, pRA-100, is maintained extrachromosomally and produces gp160 protein.

Example 38

In another construction for expressing the *env* gene, that region of HIV may be inserted into the commercially
10 available plasmid pREP4 (Invitrogen). The pREP4 plasmid is particularly useful since it contains the Epstein Barr virus origin of replication and nuclear antigen EBNA-1 coding region which produces high copy episomal replication without integration. pREP4 also contains the hygromycin marker under
15 the regulatory control of the thymidine kinase promoter and polyadenylation site. The HIV *env* coding region is placed under the regulatory control of the RSV promoter and SV40 polyadenylation site. The HIV *env* coding region was obtained as a 2.3 kb PCR fragment from HIV/3B, Genbank sequence
20 K03455. The resulting pCEP4-based plasmid, pRA-101, is maintained extrachromosomally and produces gp160 protein.

Example 39

In another construction for expressing the *gag/pol* genes, that region of HIV may be inserted into the
25 commercially available plasmid pCEP4 (Invitrogen). The pCEP4 plasmid is particularly useful since it contains the Epstein Barr virus origin of replication and nuclear antigen EBNA-1 coding region which produces high copy episomal replication without integration. pCEP4 also contains the hygromycin
30 marker under the regulatory control of the thymidine kinase promoter and polyadenylation site. The HIV *gag/pol* coding region is placed under the regulatory control of the CMV promoter and SV40 polyadenylation site. The HIV *gag/pol* coding region was obtained from HIV MN, Genbank sequence
35 MI7449, and includes the *vif* gene. The *vpr* gene is not included. The resulting pCEP4-based plasmid, pLA-100, is

- 64 -

maintained extrachromosomally and produces GAG55, r v rse transcriptase, protease and integrase proteins.

Example 40

In another construction for expressing the *gag/pol* genes, that region of HIV may be inserted into the commercially available plasmid pREP4 (Invitrogen). The pREP4 plasmid is particularly useful since it contains the Epstein Barr virus origin of replication and nuclear antigen EBNA-1 coding region which produces high copy episomal replication without integration. pREP4 also contains the hygromycin marker under the regulatory control of the thymidine kinase promoter and polyadenylation site. The HIV *gag/pol* coding region is placed under the regulatory control of the CMV promoter and SV40 polyadenylation site. The HIV *gag/pol* coding region was obtained from HIV MN, Genbank sequence MI7449, and includes the *vif* gene. The *vpr* gene is not included. The resulting pREP4-based plasmid, pLA-101, is maintained extrachromosomally and produces GAG55, reverse transcriptase, protease and integrase proteins.

Example 41

The following construction, referred to herein as pGAGPOL.rev, is useful to express HIV *gag/pol* genes.

The plasmid includes a Kanamycin resistance gene and a pBR322 origin of DNA replication. The sequences provided for transcription regulation include: a cytomegalovirus promoter; a Rous sarcoma virus enhancer; and an SV40 polyadenylation signal. The HIV-1 sequences included in pGAGPOL.rev include a sequence that encodes p17, p24, and p15 of the *gag* open reading frame; a sequence that encodes protease, a sequence that encodes reverse transcriptase which contains a small deletion and a sequence that encodes the inactive amino terminus of integrase of the *pol* open reading frame; and a sequence that encodes rev. Each of the HIV sequences are derived from HIV-1 strain HXB2.

Several safety features are included in pGAGPOL.rev. These include use of the CMV promoter and a non-retroviral poly(A) site. Furthermore, deletion of the ψ sequence limits

- 65 -

the ability to package viral RNA. In addition, multiple mutations of the reverse transcriptase yield an enzymatically inactive product. Moreover, a large deletion of integrase yields an inactive product and a Kanamycin resistance marker is used for stabilizing bacterial transformants.

Plasmid pGAGPOL.rev is constructed as follows.

Step 1. A subclone of part of the HIV-1 (HXB2) genome that is cloned into Bluescript (Stratagene) is used. The subclone of HIV-1 contains the complete 5' LTR and the rest of the HIV-1 genome to nucleotide 5795 (Genebank numbering). The HIV-1 sequences are obtained from the HXB2D plasmid (AIDS Repository).

Step 2. PCR part of gag from the open reading frame HXB2D plasmid (AIDS Repository). Cut PCR fragment with NotI and SpeI and ligate with HIV-1 subclone described above restricted with NotI and SpeI.

Step 3. PCR gag/pol junction and part of pol-encoding sequences from the HXB2D plasmid (AIDS Repository) with primers SEQ ID NO:15 and SEQ ID NO:16. Cut PCR product with ClaI and ligate together. Cut ligated fragments with BclI and SalI and ligate with plasmid from Step 2 digested with BclI and SalI.

Step 4. Cut plasmid from Step 3 with BspMI and EcoRI and religate with adapters formed by annealing linkers SEQ ID NO:17 and SEQ ID NO:18.

Step 5. Cut plasmid from Step 4 with NotI and SalI and ligate with plasmid from either 4a or 4b in description written for pENV (below). Cut also with NotI and SalI.

Step 6. Restrict plasmid from Step 5 with SalI and MluI and ligate with PCR product obtained by PCR of rev with primers SEQ ID NO:19 and SEQ ID NO:20.

Step 7. Cut plasmid from Step 6 with NotI and ligate with product obtained by PCR of the rev responsive element in the HXB2D plasmid (AIDS Repository) with primers SEQ ID NO:21 and SEQ ID NO:22.

Steps 6 and 7 are optional.

Example 42

- 66 -

The following construction, referred to herein as pENV, is useful to express HIV env genes.

The plasmid includes a Kanamycin resistance gene and a pBR322 origin of DNA replication. The sequences provided for transcription regulation include: a cytomegalovirus promoter; a Rous sarcoma virus enhancer; and an SV40 polyadenylation signal. The HIV-1 sequences included in pENV include a sequence that encodes vpu; a sequence that encodes rev; a sequence that encodes gp160; a sequence that encodes 50% of nef; a sequence that encodes vif; and, a sequence that encodes vpr with a 13 amino acid carboxy-end deletion. The vpu, rev, gp160 and nef sequences are derived from HIV-1 strain MN. The vif and vpr sequences are derived from HIV-1 strain HXB2.

Several safety features are included in pGAGPOL.rev. These include use of the CMV promoter and a non-retroviral poly(A) site. Furthermore, tat has been deleted and a 50% deletion of nef yields an "inactive" nef product. In addition, vif and vpr are placed out of normal sequence and a partial deletion of vpr further ensures an inactive vpr product.

Plasmid pENV is constructed as follows.

Step 1. Start with pUC18 digested with *Hind*III and *Eco*RI. The resulting fragment that contains the ColE1 origin of replication and the *lac*i gene should be ligated with the *Eco*RI/*Hind*III fragment from pMAMneoBlue that contains the Rous sarcoma virus enhancer. The resulting plasmid or pMAMneo-Blue from Clontech (Palo Alto, CA) can then be digested with *Hind*III and *Bgl*II. Using standard techniques, ligate with fragment containing kan gene obtained by PCR of geneblock plasmid (Pharmacia).

Step 2. If pMAMneo-Blue used as starting plasmid, digest with *Mlu*I and *Eco*RI, fill in the ends with Klenow fragment of Polymerase I and religate.

Step 3. Then, with either pMAMneo-Blue or pUC18-derived plasmid, digest with *Hind*III and ligate with the SV40 polyA site and early splicing region obtained by PCR of pCEP4

- 67 -

(Invitrogen, San Diego CA) with primers SEQ ID NO:23 and SEQ ID NO:24.

Step 4a. Digest with *Bam*HI and ligate with the CMV promoter obtained by PCR of pCEP4 (Invitrogen, San Diego CA) with primers SEQ ID NO:25 and SEQ ID NO:26.

Step 4b. Digest with *Bam*HI and ligate with the MoMLV LTR obtained by PCR with primers SEQ ID NO:27 and SEQ ID NO:28.

Step 5. Digest with *Not*I and *Mlu*I and ligate with GP160 coding region obtained by PCR of pMN-ST1 with primers SEQ ID NO:29 and SEQ ID NO:30.

Step 6. Digest with *Mlu*I and ligate with sequences that encode *vif* in its entirety and *vpr* with a 13aa carboxy-end deletion by CPR of HXB2D plasmid (AIDS Repository) with primers SEQ ID NO:31 and SEQ ID NO:32.

Example 43

An immunization system is provided which comprises:
a pharmaceutical composition comprising about 100 µg of pGAGPOL. rev in an isotonic, pharmaceutically acceptable solution; and,

a pharmaceutical preparation comprising 100 µg of pENV in an isotonic, pharmaceutically acceptable solution. In addition, the immunization system preferably comprises a pharmaceutical composition comprising about 1 ml of 0.5% bupivacaine-HCl and 0.1% methylparaben in an isotonic pharmaceutical carrier.

In such a preferred immunization system, a first set of administrations is performed in which bupivacaine and one of the two pharmaceutical compositions are administered intramuscularly to an individual, preferably into a muscle of an arm or buttock. Bupivacaine and the other of the two pharmaceutical compositions are administered intramuscularly to the individual at a different site, preferably remote from the site of the administration of the one pharmaceutical composition, preferably into a muscle of the other arm or buttock. Subsequence sets of administrations may be performed later in time, preferably 48 hours to two weeks or more later.

- 68 -

The immunization system may be used to vaccinate an individual in order to protect that individual from HIV infection or to treat an HIV infected individual with an immunotherapeutic.

5 **Example 44**

In some embodiments, the present invention relates to a method of immunizing an individual against HIV by administering a single inoculant. This inoculant includes a genetic construct that comprises at least one, preferably two,
10 more preferably more than two or a plurality of the genes of the HIV virus or all of the structural genes. However, the inoculant does not contain a complete complement of all HIV genes. If a single cell is provided with a complete complement of viral genes, it is possible that a complete
15 infectious virus can be assembled within the cell. Accordingly, a genetic construct according to the present invention is not provided with such a full complement of genes. As a safety precaution, one or more essential genes can be deleted or intentionally altered to further ensure that
20 an infectious viral particle cannot be formed.

In some embodiments of the present invention, at least portions of one, two or all HIV structural genes are provided. The structural genes of HIV consist of *gag*, *pol* and *env*. Portions of at least one of these three genes are
25 provided on a genetic construct. Accordingly, in some embodiments, at least a portion of each of *gag* and *pol* are provided on a genetic construct; in some embodiments, at least a portion of *env* is provided on a genetic construct; in some embodiments, at least a portion of *gag* is provided on a
30 genetic construct; in some embodiments at least a portion of each of *pol* and *env* are provided on a genetic construct; in some embodiments, at least a portion of each of *gag* and *env* are provided on a genetic construct; in some embodiments at least a portion of *pol* is provided on a genetic construct.
35 Optionally, the entire gene is provided. Optionally, in any of these constructs, HIV regulatory genes may also be present.

- 69 -

The HIV regulatory genes are: *vpr*, *vif*, *vpu*, *nef*, *tat* and *rev*.

Exempl 45

As used here in, the term "expression unit" is meant to refer to a nucleic acid sequence which comprises a promoter operably linked to a coding sequence operably linked to a polyadenylation signal. The coding sequence may encode one or more proteins or fragments thereof. In preferred embodiments, a expression unit is within a plasmid.

As used herein, the term "HIV expression unit" is meant to refer to a nucleic acid sequence which comprises a promoter operably linked to a coding sequence operably linked to a polyadenylation signal in which the coding sequence encodes a peptide that comprises an epitope that is identical or substantially similar to an epitope found on an HIV protein. "Substantially similar epitope" is meant to refer to an epitope that has a structure which is not identical to an epitope of an HIV protein but nonetheless invokes an cellular or humoral immune response which cross reacts to an HIV protein. In preferred embodiments, the HIV expression unit comprises a coding sequence which encodes one or more HIV proteins or fragments thereof. In preferred embodiments, an HIV expression unit is within a plasmid.

In some embodiments of the present invention, a single genetic construct is provided that has a single HIV expression unit which contains DNA sequences that encode one or more HIV proteins or fragments thereof. As used herein, the term "single HIV expression unit construct" is meant to refer to a single genetic construct that contains a single HIV expression unit. In preferred embodiments, a single HIV expression unit construct is in the form of a plasmid.

In some embodiments of the present invention, a single genetic construct is provided that has more than one HIV expression units in which each contain DNA sequences that encode one or more HIV proteins or fragments thereof. As used herein, the term "multiple HIV expression unit genetic construct" is meant to refer to a single plasmid that contains more than one HIV expression units. In preferred embodiments,

- 70 -

a multiple HIV expression unit construct is in the form of a plasmid.

In some embodiments of the present invention, a single genetic construct is provided that has two HIV expression units in which each contain DNA sequences that encode one or more HIV proteins or fragments thereof. As used herein, the term "two HIV expression unit genetic construct" is meant to refer to a single plasmid that contains two HIV expression units, i.e. a multiple HIV expression unit genetic construct that contains two HIV expression unit genetic expression units. In a two HIV expression unit genetic construct, it is preferred that one HIV expression unit operates in the opposite direction of the other HIV expression unit. In preferred embodiments, a two HIV expression unit construct is in the form of a plasmid.

In some embodiments of the present invention, an HIV genetic vaccine is provided which contains a single genetic construct. The single genetic construct may be a single HIV expression unit genetic construct, a two HIV expression unit genetic construct or a multiple HIV expression unit genetic construct which contains more than two HIV expression units.

In some embodiments of the present invention, an HIV genetic vaccine is provided which contains more than one genetic construct in a single inoculant.

In some embodiments of the present invention, an HIV genetic vaccine is provided which contains more than one genetic construct in more than one inoculant. As used herein, the term "multiple inoculant" is meant to refer to a genetic vaccine which comprises more than one genetic construct, each of which is administered separately. In some embodiments of the present invention, an HIV genetic vaccine is provided which contains two genetic constructs. Each genetic construct may be, independently, a single HIV expression unit genetic construct, a two HIV expression unit genetic construct or a multiple HIV expression unit genetic construct which contains more than two HIV expression units. In some embodiments, both genetic constructs are single HIV expression unit genetic

- 71 -

constructs. In some embodiments, both genetic constructs are two HIV expression unit genetic constructs. In some embodiments, both genetic constructs are multiple HIV expression unit genetic constructs. In some embodiments, one
5 genetic construct is a single HIV expression unit genetic construct and the other is a two HIV expression unit genetic construct. One having ordinary skill in the art can readily recognize and appreciate the many variations depending upon the number of genetic constructs used in a genetic vaccine and
10 the number of HIV expression units that may be present on each genetic construct.

It is preferred that the genetic constructs of the present invention do not contain certain HIV sequences, particularly, those which play a role in the HIV genome
15 integrating into the chromosomal material of the cell into which it is introduced. It is preferred that the genetic constructs of the present invention do not contain LTRs from HIV. Similarly, it is preferred that the genetic constructs of the present invention do not contain a *psi* site from HIV.
20 Further, it is preferred that the reverse transcriptase gene is deleted and the integrase gene is deleted. Deletions include deletion of only some of the codons or replacing some of the codons in order to essentially delete the gene. For example, the initiation codon may be deleted or changed or
25 shifted out of frame to result in a nucleotide sequence that encodes an incomplete and non-functioning.

It is also preferred that the genetic constructs of the present invention do not contain a transcribable *tat* gene from HIV. The *tat* gene, which overlaps the *rev* gene may be
30 completely deleted by substituting the codons that encode *rev* with other codons that encode the same amino acid for *rev* but which does not encode the required *tat* amino acid in the reading frame in which *tat* is encoded. Alternatively, only some of the codons are switched to either change, i.e.
35 essentially delete, the initiation codon for *tat* and/or change, i.e. essentially delete, sufficient codons to result

- 72 -

in a nucleotide sequence that encodes an incomplete and non-functional tat.

It is preferred that a genetic construct comprises coding sequences that encode peptides which have at least an epitope identical to or substantially similar to an epitope from HIV gag, pol, env or rev proteins. It is more preferred that a genetic construct comprises coding sequences that encode at least one of HIV gag, pol, env or rev proteins or fragments thereof. It is preferred that a genetic construct comprises coding sequences that encode peptides which have more than one epitopes identical to or substantially similar to an epitope from HIV gag, pol, env or rev proteins. It is more preferred that a genetic construct comprises coding sequences that encode more than one of HIV gag, pol, env or rev proteins or fragments thereof.

In some embodiments, a genetic construct comprises coding sequences that encode peptides which have at least an epitope identical to or substantially similar to an epitope from HIV vif, vpr, vpu or nef proteins. In some embodiments, a genetic construct comprises coding sequences that encode at least one of HIV vif, vpr, vpu or nef proteins or fragments thereof.

A single HIV expression unit genetic construct may comprise coding regions for one or more peptides which share at least one epitope with an HIV protein or fragment thereof in a single expression unit under the regulatory control of single promoter and polyadenylation signal. It is preferred that genetic constructs encode more than one HIV protein or fragment thereof. The promoter may be any promoter functional in a human cell. It is preferred that the promoter is an SV40 promoter or a CMV promoter, preferably a CMV immediate early promoter. The polyadenylation signal may be any polyadenylation signal functional in a human cell. It is preferred that the polyadenylation signal is an SV40 polyadenylation signal, preferably the SV40 minor polyadenylation signal. If more than one coding region is provided in a single expression unit, they may be immediately

- 73 -

adjacent to each other or separated by non-coding regions. In order to be properly expressed, a coding region must have an initiation codon and a termination codon.

A two HIV expression unit genetic construct may
5 comprise coding regions for one or more peptides which share at least one epitope with an HIV protein or fragment thereof on each of the two expression units. Each expression unit is under the regulatory control of single promoter and polyadenylation signal. In some embodiments, it is preferred
10 that genetic constructs encode more than one HIV protein or fragment thereof. In some embodiments, it is preferred that nucleotide sequences encoding *gag* and *pol* are present on one expression unit and nucleotide sequences encoding *env* and *rev* are present on the other. The promoter may be any promoter
15 functional in a human cell. It is preferred that the promoter is an SV40 promoter or a CMV promoter, preferably a immediate early CMV promoter. The polyadenylation signal may be any polyadenylation signal functional in a human cell. It is preferred that the polyadenylation signal is an SV40
20 polyadenylation signal, preferably the SV40 minor polyadenylation signal. If more than one coding region is provided in a expression unit, they may be immediately adjacent to each other or separated by non-coding regions. In order to be properly expressed, a coding region must have
25 an initiation codon and a termination codon.

According to some embodiments of the present invention, the MHC Class II crossreactive epitope in *env* is deleted and replaced with the analogous region from HIV II.

When a genetic construct contains *gag* and/or *pol*,
30 it is generally important that *rev* is also present. In addition to *rev*, a *rev* response element may be provided with *gag* and *pol* for increased expression of those genes.

When genetic constructs are produced that it is preferred that the *env* gene used in plasmid 1 is derived from
35 MN or MN-like isolates including clinical isolates resembling MN, preferably non-syncytial inducing clinical isolates,

- 74 -

preferably those that are macrophage tropic from early stage clinical isolates.

Multiple proteins may be produced from a single expression unit by alternative splicing. Splicing signals are provided to allow alternative splicing which produces different messages encoding different proteins.

Example 46

Figure 8 shows four backbones, A, B, C and D. Figure 9 shows 4 inserts, 1, 2, 3 and 4. Insert 1 supports expression of *gag* and *pol*; the *rev* response element was cloned in a manner to conserve the HIV splice acceptor. Insert 2 is similar to insert 1 as it too supports expression of *gag* and *pol* except the *rev* response element was cloned without conserving the HIV splice acceptor. Insert 3 supports expression of *gag* and *pol*, includes a deletion of the integrase gene and does not include the presence of the *cis* acting *rev* response element. Insert 4 supports expression of *rev*, *vpu* and *env*. The *env* may have the MHC class II cross reactive epitope altered to eliminate crossreactivity and the V3 loop may be altered to eliminate the possibility of syncytia formation.

In some embodiments, backbone A is used with insert 1. Such constructs optionally contain the SV40 origin of replication. Plasmid pA1ori+ is backbone A with insert 1 and the SV40 origin of replication. Plasmid pA1ori- is backbone A with insert 1 without the SV40 origin of replication. Additionally, either pA1ori+ or pA1ori- may include integrase yielding pA1ori+int+ and pA1ori-int+, respectively. Plasmids pA1ori+, pA1ori-, pA1ori+int+ and pA1ori-int+ may be further modified by functionally deleting the reverse transcriptase (RT) gene yielding pA1ori+RT-, pA1ori-RT-, pA1ori+int+RT- and pA1ori-int+RT-, respectively.

In some embodiments, backbone A is used with insert 2. Such constructs optionally contain the SV40 origin of replication. Plasmid pA2ori+ is backbone A with insert 2 and the SV40 origin of replication. Plasmid pA2ori- is backbone A with insert 1 without the SV40 origin of replication.

- 75 -

Additionally, either pA2ori+ or pA2ori- may include integrase yielding pA2ori+int+ and pA2ori-int+, respectively. Plasmids pA2ori+, pA2ori-, pA2ori+int+ and pA2ori-int+ may be further modified by functionally deleting the reverse transcriptase (RT) gene yielding pA2ori+RT-, pA2ori-RT-, pA2ori+int+RT- and pA2ori-int+RT-, respectively.

In some embodiments, backbone B is used with insert 1. Such constructs optionally the SV40 origin of replication. Plasmid pB1ori+ is backbone B with insert 1 and the SV40 origin of replication. Plasmid pB1ori- is backbone B with insert 1 without the SV40 origin of replication. Additionally, either pB1ori+ or pB1ori- may include integrase yielding pB1ori+int+ and pB1ori-int+, respectively. Plasmids pB1ori+, pB1ori-, pB1ori+int+ and pB1ori-int+ may be further modified by functionally deleting the reverse transcriptase (RT) gene yielding pB1ori+RT-, pB1ori-RT-, pB1ori+int+RT- and pB1ori-int+RT-, respectively.

In some embodiments, backbone B is used with insert 2. Such constructs optionally the SV40 origin of replication. Plasmid pB2ori+ is backbone B with insert 2 and the SV40 origin of replication. Plasmid pB2ori- is backbone B with insert 1 without the SV40 origin of replication. Additionally, either pB2ori+ or pB2ori- may include integrase yielding pB2ori+int+ and pB2ori-int+, respectively. Plasmids pB2ori+, pB2ori-, pB2ori+int+ and pB2ori-int+ may be further modified by functionally deleting the reverse transcriptase (RT) gene yielding pB2ori+RT-, pB2ori-RT-, pB2ori+int+RT- and pB2ori-int+RT-, respectively.

In some embodiments, backbone A minus rev is used with insert 3. Such constructs optionally the SV40 origin of replication. Plasmid pA/r-3ori+ is backbone A with insert 2 and the SV40 origin of replication. Plasmid pA/r-3ori- is backbone A minus rev with insert 3 without the SV40 origin of replication. Additionally, either pA/r-3ori+ or pA/r-3ori- may include integrase yielding pA/r-3ori+int+ and pA/r-3ori-int+, respectively. Plasmids pA/r-3ori+, pA/r-3ori-, pA/r-3ori+int+ and pA/r-3ori-int+ may be further modified by

- 76 -

functionally deleting the reverse transcriptase (RT) gene yielding pA/r-3 ri+RT-, pA/r-3ori-RT-, pA/r-3ori+int+RT- and pA/r-3ori-int+RT-, respectively.

In some embodiments, backbone C is used with insert

- 5 1. Such constructs optionally the SV40 origin of replication. Plasmid pC1ori+ is backbone C with insert 1 and the SV40 origin of replication. Plasmid pC1ori- is backbone C with insert 1 without the SV40 origin of replication. Additionally, either pC1ori+ or pC1ori- may include integrase
10 yielding pC1ori+int+ and pC1ori-int+, respectively. Plasmids pC1ori+, pC1ori-, pC1ori+int+ and pC1ori-int+ may be further modified by functionally deleting the reverse transcriptase (RT) gene yielding pC1ori+RT-, pC1ori-RT-, pC1ori+int+RT- and pC1ori-int+RT-, respectively.

- 15 In some embodiments, backbone C is used with insert 2. Such constructs optionally the SV40 origin of replication. Plasmid pC2ori+ is backbone C with insert 2 and the SV40 origin of replication. Plasmid pC2ori- is backbone C with insert 2 without the SV40 origin of replication.
20 Additionally, either pC2ori+ or pC2ori- may include integrase yielding pC2ori+int+ and pC2ori-int+, respectively. Plasmids pC2ori+, pC2ori-, pC2ori+int+ and pC2ori-int+ may be further modified by functionally deleting the reverse transcriptase (RT) gene yielding pC2ori+RT-, pC2ori-RT-, pC2ori+int+RT- and
25 pC2ori-int+RT-, respectively.

- In some embodiments, backbone C is used with insert 3. Such constructs optionally the SV40 origin of replication. Plasmid pC3ori+ is backbone C with insert 3 and the SV40 origin of replication. Plasmid pC3ori- is backbone C with
30 insert 3 without the SV40 origin of replication. Additionally, either pC3ori+ or pC3ori- may include integrase yielding pC3ori+int+ and pC3ori-int+, respectively. Plasmids pC3ori+, pC3ori-, pC3ori+int+ and pC3ori-int+ may be further modified by functionally deleting the reverse transcriptase
35 (RT) gene yielding pC3ori+RT-, pC3ori-RT-, pC3ori+int+RT- and pC3ori-int+RT-, respectively.

- 77 -

In some embodiments, backbone D is used with insert 4. Such constructs optionally the SV40 origin of replication. Plasmid pD4ori+ is backbone D with insert 4 and the SV40 origin of replication. Plasmid pD4ori- is backbone D with insert 4 without the SV40 origin of replication.

Example 47

In some embodiments, a single expression unit/single inoculant genetic vaccine is provided which comprises a genetic construct that includes a coding sequence which encodes a peptide that has at least one epitope which is identical to or substantially similar to epitopes of HIV proteins. The coding sequence is under the regulatory control of the CMV immediate early promoter and the SV40 minor polyadenylation signal.

In some embodiments, a single expression unit/single inoculant genetic vaccine is provided which comprises a genetic construct that includes a coding sequence which encodes at least one HIV protein or a fragment thereof. The coding sequence is under the regulatory control of the CMV immediate early promoter and the SV40 minor polyadenylation signal. The HIV protein is selected from the group consisting of *gag*, *pol*, *env* and *rev*. In some embodiments it is preferred that the genetic vaccine is provided which comprises a genetic construct that includes a coding sequence which encodes at least two HIV proteins or a fragments thereof selected from the group consisting of *gag*, *pol*, *env* and *rev* or fragments thereof. In some embodiments, it is preferred that the genetic vaccine is provided which comprises a genetic construct that includes a coding sequence which encodes at least three HIV proteins or a fragments thereof selected from the group consisting of *gag*, *pol*, *env* and *rev* or fragments thereof. In some embodiments, it is preferred that the genetic vaccine is provided which comprises a genetic construct that includes a coding sequence which encodes *gag*, *pol*, *env* and *rev* or fragments thereof.

In some embodiments, a dual expression unit/single inoculant genetic vaccine is provided which comprises a

- 78 -

genetic construct that includes two expression units each of which comprises a coding sequence which encodes a peptide that has at least one epitope which is an identical to or substantially similar to epitopes of HIV proteins. The coding
5 sequence is under the regulatory control of the CMV immediate early promoter and the SV40 minor polyadenylation signal. The two expression units are encoded in opposite directions of each other.

In some embodiments, a dual expression unit/single
10 inoculant genetic vaccine is provided which comprises a genetic construct that includes two expression units each of which comprises a coding sequence which encodes at least one HIV protein or a fragment thereof. Each expression unit
15 comprises a coding sequence that is under the regulatory control of the CMV immediate early promoter and the SV40 minor polyadenylation signal. The HIV protein is selected from the group consisting of *gag*, *pol*, *env* and *rev*. In some
20 embodiments it is preferred that the genetic vaccine is provided which comprises a genetic construct that includes two expression units, at least one of which comprises a coding
25 which encodes at least two HIV proteins or a fragments thereof selected from the group consisting of *gag*, *pol*, *env* and *rev* or fragments thereof and the other comprises at least one HIV
30 proteins or a fragments thereof selected from the group consisting of *gag*, *pol*, *env* and *rev* or fragments thereof. In some embodiments, it is preferred that the genetic vaccine is
35 provided which comprises a genetic construct that includes two expression units, at least one of which comprises a coding sequence which encodes at least three HIV proteins or a
fragments thereof selected from the group consisting of *gag*, *pol*, *env* and *rev* or fragments thereof and the other comprises at least one HIV proteins or a fragments thereof selected from the group consisting of *gag*, *pol*, *env* and *rev* or fragments thereof. In some embodiments, it is preferred that the
genetic vaccine is provided which comprises a genetic
construct that comprises two expression units and includes a

- 79 -

coding sequence which encodes *gag*, *pol*, *env* and *rev* or fragments thereof.

Example 48

A genetic construct, plasmid pCMN160Δ16 was made for use in an anti-HIV pharmaceutical kit or pharmaceutical composition. pCMN160Δ16 was constructed as follows:

Step 1: Primers SEQ ID NO:35 and SEQ ID NO:34 were used a PCR fragment from HIV/MN genomic DNA.

Step 2: Primers SEQ ID NO:33 and SEQ ID NO:36 were used a PCR fragment from HIV/MN genomic DNA.

Step 3: Primers SEQ ID NO:35 and SEQ ID NO:36 were combined with 2 μl of reaction material from Steps 1 and 2.

Step 4: Reaction product from Step 3 was cut with *Not*I and *Mlu*I and inserted into Backbone A described in Example 46 cut with *Not*I and *Mlu*I.

Plasmid pCMN160Δ16 is thereby formed which contains as an insert to Backbone A a coding region which encodes the MN strain ENV Protein with the *rev* region and half of *nef* having HLA-DB region changes to HIV-2.

Example 49

The plasmid pGAGPOL.rev2 was made as follows. First the backbone was made. Then an insert with HIV *gag* and *pol* was generated and inserted into the backbone.

The backbone was prepared as follows.

Step 1. Digest pMAMneo (Clonetech) with *Bgl*II. Fill-in with Klenow fragment of Polymerase I. Cut with *Hind*III. Gel purify 1763bp fragment.

Step 2. Amplify Kan^R gene from plasmid pJ40kan⁺ (Kanamycin resistance gene obtained from Pharmacia Inc. cloned into pJ40 obtained as a gift from the Imperial Cancer Research Fund UK; pJ40 was originally constructed and reported by Morgenstern, J.P. and H. Land, *Nucl. Acids Res.* 18(4):1068, which is incorporated herein by reference) with oligos SEQ ID NO:37 and SEQ ID NO:38. Blunt off PCR product. Cut with *Hind*III. Gel purify PCR fragment.

Step 3. Ligate the vector backbone generated from pMAMneo and described in step #1 with the PCR product encoding

- 80 -

the Kan^R gen and described in step #2. Isolate plasmid containing the Kan^R gene and the bacterial origin of replication.

Step 4. Digest resulting plasmid with *Mlu*I, fill-in with Klenow fragment of DNA polymerase I. Ligate with *Sac*II linker (New England Biolabs).

Step 5. Digest plasmid obtained in step 4 with *Ase*I and *Ssp*I.

Step 6. PCR part of the Kan^R gene from the plasmid described in step 3 using primers SEQ ID NO:39 and SEQ ID NO:40. Cut PCR product with *Ssp*I and *Ase*I.

Step 7. Ligate largest fragment obtained in step 5 with PCR product obtained in step 6.

Step 8. Cut ligation product/plasmid obtained in step 7 with *Hind*III. Blunt off with the Klenow fragment of DNA polymerase I.

Step 9. Cut pCEP4 (Invitrogen) with *Sal*I to release a DNA fragment containing the CMV promoter, polylinker, and SV40 poly A site. Purify this fragment and blunt-off with the Klenow fragment of DNA Polymerase I.

Step 10. Ligate the plasmid obtained in step 8 and the fragment obtained in step 9. Isolate plasmid containing the bacterial origin of replication, the Kan^R gene, the RSV enhancer, the CMV promoter, polylinker, and the SV40 poly A site.

Step 11. Cut plasmid obtained in step 10 with *Bam*HI and *Nhe*I.

Step 12. Anneal oligonucleotides SEQ ID NO:41 and SEQ ID NO:42.

Step 13. Ligate the plasmid obtained in step 10 with the annealed oligonucleotides obtained in step 12. Isolate plasmid containing the adapter contained in step 12.

Step 14. Digest plasmid obtained in step 13 with *Sal*I and *Mlu*I.

Step 15. PCR amplify the rev open reading frame using BBG35 (RD Systems Inc. Minneapolis, MN; which contains the coding region for rev from HIV strain HX3B in pUC19) as

- 81 -

a template and primers SEQ ID NO:43 and SEQ ID NO:44. Digest the PCR product with *SalI* and *MluI*.

Step 16. Ligate the plasmid obtained in step 14 with the PCR product produced in step 15. Isolate plasmid 5 containing the rev coding region.

Preparation of gag/pol insert.

Step 1. A subclone of part of the HIV-1 (HXB2) genome that was cloned into Bluescript (Stratagene). The subclone of HIV-1 contains the complete 5'LTR and the rest of 10 the HIV-1 genome to nucleotide 5795 (Genbank numbering) cloned into the *XbaI* and *SalI* sites of Bluescript. The HIV-1 sequences are obtained from the HXB2D plasmid (AIDS Repository).

Step 2. PCR part of the gag coding region from the 15 open reading frame of the plasmid described in step 1 (the subclone of part of the HIV-1 HXB2 genome that is cloned into Bluescript) using primers SEQ ID NO:45 and SEQ ID NO:46:

Step 3. Digest plasmid described in step 1 (the subclone of part of the HIV-1 HXB2 genome that is cloned into 20 Bluescript) with *EcoRI*. Purify the plasmid that contains the pBluescript backbone, the 5' HIV-1 LTR, the gag coding region and part of the pol coding region and religate.

Step 4. Cut the plasmid obtained in step 3 with *NotI* and *SpeI* and ligate with the PCR fragment described in 25 Step 2 after it is digested with *NotI* and *SpeI*. Isolate plasmid that contain the PCR fragment instead of the original *NotI/SpeI* fragment which contains the 5' HIV-1 LTR.

Step 5. Digest the plasmid obtained in step 4 with *EcoRI* and *SalI*.

30 Step 6. Anneal oligonucleotides SEQ ID NO:47 and SEQ ID NO:48.

Step 7. Ligate the plasmid obtained in step 5 with the adapter obtained in step 6. Isolate plasmid containing the adapter cloned into the *EcoRI/SalI* sites.

35 Step 8. Digest the plasmid obtained in step 7 with *NdeI* and *EcoRI*.

- 82 -

Step 9. PCR amplify the Rev Response Element (RRE) from a plasmid containing the RRE sequence from HIV-1 strain HXB2 using primers SEQ ID NO:49 and SEQ ID NO:50. Digest the PCR product with *NdeI* and *EcoRI*.

5 Step 10. Ligate the plasmid obtained in step 8 with the PCR product obtained in step 9. Isolate plasmid containing the insert with the RRE sequence.

 Step 11. Digest the plasmid obtained in step 10 with *NotI* and *Sall* and isolate the fragment containing the *gag* coding region, the modified *pol* coding region, and the RRE sequence.

 Step 12. Digest the plasmid obtained in step 16 of the protocol for preparing the backbone which is described above with *NotI* and *Sall*.

15 Step 13. Ligate the plasmid obtained in step 12 with the insert obtained in step 11. Isolate plasmids that contain the insert containing the *gag* coding region, the modified *pol* coding region, and the RRE sequence.

 Step 14. Digest plasmid obtained in step 13 with *XbaI* and *NheI*, Blunt-off ends and religate. Isolate the plasmid that is lacking the *KpnI* site that is present between the *XbaI* and *NheI* sites in the plasmid obtained in step 13.

 Step 15. Digest the plasmid obtained in step 14 with *KpnI* and isolate the largest fragment.

25 Step 16. Anneal oligonucleotides SEQ ID NO:51 and SEQ ID NO:52.

 Step 17. Ligate the purified plasmid fragment obtained in step 15 with the adapter obtained in step 16. Isolate plasmid containing the adapter inserted at the *KpnI* site of the plasmid obtained in step 15.

Example 50 Genetic Immunization with Genes for Regulatory Proteins

 Part of the difficulty of combatting HIV arises from the extraordinary variability of the virus and its ability to quickly mutate into new forms. Not only is there substantial protein sequence variation among HIV isolates found in the human population as a whole, but the virus mutates so quickly

- 83 -

that every HIV-infected individual actually harbors a number of related HIV microvariants. Such HIV isolates exhibit differences in replication efficiency, tropism, susceptibility to neutralization, and drug resistance. As drug-resistant mutants appear, the benefits of drug therapy fade. With AZT, drug resistance typically arises within the first year of therapy. This constant generation of escape mutants may play a part in the ability of HIV to finally overwhelm host defenses after a long period in which the virus appears to be held in check.

This mutational drift has been reported in various regions of the gp120 envelope glycoprotein, including the principal neutralizing domain of the V3 loop, and in the HIV core proteins as well. HIV regulatory proteins are much more highly conserved than the structural proteins and also exhibit less mutational drift over time. Regulatory proteins therefore present attractive targets for antiviral attack.

HIV exhibits a remarkable temporal regulation of expression of regulatory vs structural proteins. In the early phase of viral replication, mRNAs encoding the regulatory proteins Tat, Rev and Nef predominate, whereas in the late phase, there is greater expression of mRNAs encoding structural proteins, including Gag, Pol, and Env precursors, and many accessory proteins. This shift from early to late phase is triggered when the Rev protein reaches a particular level. The predominance of Tat, Rev and Nef early in the viral replication cycle also makes these proteins favorable targets for antiviral attack. This is especially true for tat and rev, which play absolutely essential roles in transcriptional and post-translational regulation of HIV gene expression, and predominate early in the viral replication cycle, before transcription of viral structural proteins and production of infectious viral particles.

In contrast to tat and rev, which clearly play essential roles in HIV replication, other regulatory proteins such as nef, vpr, vif, and vpu are sometimes referred to as "accessory" proteins. Their functions are less well

- 84 -

understood, and the degree to which viral replication is attenuated by loss of a particular function varies considerably and may depend on the host cell being infected. Nevertheless, the strong conservation of such functions among
5 widely diverse HIV isolates, as well as other primate immunodeficiency viruses, suggests the importance of these "accessory" functions in the natural infection process. (See in general, Terwilliger, E.F., (1992) *AIDS Research Reviews* 2:3-27, W.C. Koff, F. Wong-Staal, and R.C. Kennedy, eds. (New
10 York:Marcel Dekker, Inc.). In fact, primate recombinant viruses deleted in either *vpr*, *nef* or *vif* are non-pathogenic *in vivo*, further demonstrating the importance of these accessory genes in the life cycle of the virus.

There is some evidence that higher level, more
15 protective immune responses against HIV could be achieved by presenting a select few regulatory and/or enzymatic proteins, rather than the entire complement of HIV genes. Accordingly, a focused immunization strategy may desirably involve genetic immunization using coding sequences for one or more
20 regulatory, non-structural HIV proteins, including *tat*, *rev*, *vpr*, *nef*, *vpu* or *vif*. Only *vpr* has been found to be associated with viral particles, whereas other regulatory proteins, including *tat*, *rev*, *nef*, *vif* and *vpu*, are not virion associated.

25 In some embodiments of genetic immunization against HIV using regulatory genes, the one or more of *tat*, *rev*, *nef*, *vif* and *vpu* genes are inserted into backbone A which is described in Example 46. It is preferred that *tat* and/or *rev* is used. In some embodiments, *tat* or *rev* are inserted into
30 backbone A which is described in Example 46. In some embodiments, Next in descending order of desirability as targets are *nef*, *vpr*, *vif*, and *vpu*. Preferably, more than one regulatory gene will be employed, including *tat* and *rev*; *tat*, *rev*, and *nef*; *tat*, *rev*, *nef*, and *vpr*; *tat*, *rev*, *nef*, *vpr*,
35 and *vif*; *tat*, *rev*, *nef*, *vpr*, *vif*, and *vpu*; as well as combinations thereof; and, optionally, such additional regulatory genes as *tev*.

- 85 -

The Tat protein is a transactivator of LTR-directed gene expression. It is absolutely essential for HIV replication. Tat is produced early in the viral replication cycle and functional Tat is required for expression of Gag, Pol, Env and Vpr. The predominant form of Tat is an 86-amino acid protein derived from two exon mRNAs. The amino-terminal 58 amino acids are sufficient for transactivation, although with reduced activity. Tat acts on a *cis*-acting sequence termed *tar*, to produce a dramatic increase in LTR-driven gene expression. Tat may act in part through increased RNA synthesis and in part by increasing the amount of protein synthesized per RNA transcript. Until recently, Tat was thought to act only on the HIV-1 LTR. However, Tat-activated expression from the JC virus late promoter has also been reported. Tat may also stimulate cell proliferation as an exogenous factor, and may play a contributory role in promoting the growth of Kaposi's Sarcoma in HIV-infected individuals. Because of such potentially detrimental effects in both HIV-infected and -noninfected individuals, preferred tat constructs employed for genetic immunization are modified to express only non-functional Tat. Mutations capable of inactivating Tat or Rev can in addition act as transdominant mutations, thereby potentially inactivating any functional Tat being produced in an HIV-infected individual.

Rev is a second regulatory protein of HIV that is essential for viral replication. It is a 19 kD (116 amino acid) protein which is expressed from two coding exons found in a variety of multiply spliced mRNAs. Two distinct domains have been identified, a basic region involved in binding to RRE (Rev-response-element)containing transcripts and an "activation" domain that induces nuclear exports of such transcripts as a result of binding. In the course of natural viral infection, Rev is required for expression of the HIV structural proteins Gag, Pol, and Env, as well as Vpr.

Vpr is a 15 kD protein (96 amino acids) in most HIV-1 strains, although the Vpr open reading frame is extensively truncated in many viral strains extensively passaged in cell

- 86 -

cultur . Th vpr open reading frame is also present in HIV-2 and m st SIV isolates. Vpr is the first retroviral regulatory prot in found to be associated with HIV viral particles. Its presence in the HIV virion suggests it may serve a function
5 at some early point in the viral replication cycle. Vpr accelerates HIV replication, especially early in infection. Vpr increases the level of expression of reporter genes linked to the HIV LTR by about three fold. Moreover, Vpr and Tat appear to act synergistically with respect to LTR-linked
10 genes. Vpr can be isolated from the serum of HIV-infected individuals and appears to increase the ability of the virus to infect new cells. Vpr has also been found to inhibit cell proliferations and to induce cell differentiation (Levy, D.N. et al., *Cell* (1993) 72:1-20), a finding that may be
15 significant in view of reports that primary monocyte/macrophages are infectible in vitro only while undergoing differentiation (Schuitemaker, H. et al., (1992) *J. Clin. Invest.* 89:1154-1160. Even cells that are unable to support HIV replication may be disregulated by the effects of
20 Vpr. For example, Vpr may be responsible for the muscle wasting frequently observed in AIDS patients. Because of the potentially detrimental activity of Vpr, genetic immunization should preferably be carried out with a modified vpr construct which will express a non-functional Vpr protein.

25 Nef (also called 3' orf in older literature) is a 25-27kD protein. It has been suggested that Nef may be involved in the downregulation of CD4+ T lymphocytes. In addition, Nef may play a role in cell signaling. Nef appears to be important for the establishment of HIV infection in
30 vivo. Nef-specific CTLs are believed to be important in controlling HIV infection in vivo.

Vif is a 23 kD cytoplasmic protein designated "viral infectivity factor". Although Vif-defective mutant viruses are not compromised with respect to cell-to-cell transmission,
35 they exhibit a profound decrease in ability to infect many CD4+ cell lines. Without Vif, there is decreased budding of virus, and decreased infectivity. In primate studies, Vif

- 87 -

deletion mutants exhibit a severely diminished ability to establish infection *in vivo*. These studies support a clinical role for Vif in virus replication in the host.

Vpu is a 15-20 kD (81 amino acid) protein. Although
5 Vpu(+) and Vpu(-) viruses produce the same amount of viral protein, the latter exhibit increased intracellular accumulation of viral proteins together with decreased extracellular virus. This suggests that Vpu may be involved in the assembly and/or release of viral particles.

10 Simple retroviruses, such as murine and avian viruses, lack proteins analogous to the HIV-1, HIV-2, and SIV regulatory proteins. In such animals retroviral infection tends to be self-limiting, with clearance of virus and decreased pathogenicity. Similarly, HTLV-1, which includes
15 only Tax (which acts much like Tat and also exhibits vpr-like activity) and Rex (which acts much like Rev) is cleared in many individuals. Genetic immunization with regulatory genes is considered relevant not only for HIV, but also for viruses such as HBV (X gene product) and HCV, and HTLV-1 (Tax) and
20 (Rex). In all of these viruses the regulatory genes are believed to play a critical role in the virus life cycle and the establishment of infection.

Example 51 Construction of HIV-1 Regulatory Plasmid, pREG

The pREG plasmid is constructed in a stepwise
25 fashion, and each intermediate can be tested for protein expression before construction is continued. An expression vector supporting the expression of tat and rev is constructed via two steps. First, an amplification product containing a 5' NheI site, the HIV-1 major splice donor site, the majority
30 of the tat coding region, the region encoding the amino terminal region of the rev protein and an AvaII site is amplified from a synthetic template. This synthetic template is generated using the published sequences of HXB2 strain of HIV-1 obtained from the GenBank Database, and is altered to
35 mutate the cysteine residues at positions 22 and 30 of the tat protein. These mutations have been shown to render tat non-

- 88 -

functional (Kuppuswamy, et al. (1989) *Nucleic Acids Research* 17(9): 3551-3561).

The PCR product is ligated into a vector that is digested with *NheI* and *AvaII* and which contains a kanamycin resistance gene and a pBR322 origin of replication. In addition, this plasmid contains a cytomegalovirus promoter, a Rous sarcoma virus enhancer, the rev coding region and a SV40 polyadenylation signal. The rev sequence present in the plasmid is derived from the proviral clone of HIV-1 III₂. This will generate an expression vector containing a complete, but mutated, *tat* coding region and a complete rev coding region.

The subsequent step is performed to generate a PCR product containing an *AvaII* site at its 5' end, a mutation at amino acid position 81 of rev, approximately 30% of the rev coding region, approximately 30% of the *nef* coding region, and a *MluI* site at the 3' end. The amino acid change at position 81 has been shown to eliminate rev function, and therefore, the resulting plasmid will lead to production of non-functional rev protein (Bogard, H. and Greene, W.C. (1993) *J. Virol.* 67(5):2496-2502). It is assumed that the major deletion of the *nef* coding region will result in production of a non-functional *nef* protein. The 5' *AvaII* site and the mutation at amino acid position 81 of the rev protein are introduced on the 5' PCR primer which is complementary to the coding region of rev containing both the *AvaII* site and the nucleotide encoding amino acid 81. A stop codon causing termination of *Nef* at amino acid position 63 and the 3' coding cloning site, *MluI*, will be introduced by the 3' PCR primer. The template for this PCR amplification is a plasmid or synthetic template containing the rev and *nef* coding regions from the MN strain of HIV-1. The resulting PCR product will be digested with *AvaII* and *MluI*, and used to replace the smaller *AvaII*-*MluI* fragment which results after digestion of the *tat*-rev plasmid described in the preceding paragraph with *AvaII* and *MluI*.

- 89 -

Opti nally, vpr can b added to this plasmid in one of two sites. In one approach, vpr can be amplified using a 5' PCR primer c ntaining *MluI* site upstream of sequences which span the vpr translational start codon and a 3' PCR primer
5 complementary to the vpr stop codon and sequences that flank it which also contain a *MluI* cloning site. Sequences upstream of the start codon contain a splice acceptor. The PCR product can be digested with *MluI* and inserted into the *tat rev nef* plasmid described above after its digestion with *MluI*.

10 Alternatively, the vpr amplification can be performed in analogous manner, however, the PCR primers would contain restriction sites compatible with cloning into another vector so that it is expressed under the control of a second eukaryotic promoter. The cassette derived from this plasmid,
15 containing the second promoter followed by the vpr coding region, followed by the a polyA sequence, could be released by digestion with restriction enzymes that flank the cassette, but do not cut within it. The resulting DNA fragment would be cloned into a unique site of the *tat, rev, vpr* plasmid that
20 falls outside of the region necessary for the expression of *tat rev vpr*. In this way, a plasmid having two expression units is formed.

Example 52 Construction of HCV and HTLV-1 Plasmids

A similar approach can be used to generate a plasmid
25 expressing HTLV-1 or HCV encoded proteins having enzymatic functions required for the viral life cycle and/or for the regulatory proteins of these viruses. For HTLV-1, a plasmid encoding the regulatory protein, TAX, is generated using the a plasmid backbone and a cloning strategy similar to those
30 described above. Such HCV genes that encode enzymatic proteins include the RNA-dependent RNA-polymerase, a protein having helicase/protease function. The sequences necessary are published and available through GenBank. The viral organization of HTLV-1 and HCV are published in Cann, A.J. and
35 Chen, I.S.Y. *Virology* 2nd Edition, edited by B.N. Fiddr, Raven Press, Ltd., New York, 1990 and Bradley, D.W. *Transfusion Medicine Reviews*, 1(2):93-102, 1992, respectively.

- 90 -

Example 53 Genetic Immunization with Enzymatic Genes

Genetic immunization with genes encoding proteins with enzymatic functions, such as the HIV pol gene can also be an important antiviral strategy since enzymes such as Pol are necessary for the production of live virus. Without polymerase or any of its component functions, HIV is non-pathogenic and non-infectious. Similarly, the enzymatic genes of other viruses, such as the HBV polymerase, are attractive targets for genetic immunization. See, e.g., Radziwill et al., Mutational Analysis of the Hepatitis B Virus P Gene Product: Domain Structure and RNase H Activity, J. Virol. 64 (2): 613-620 (1990).

One reason for the attractiveness of viral enzymes as an immunological target is the limited ability of such enzymes to mutate their amino acid sequence and still maintain their enzymatic functions. For example, with HIV-1, Pol exhibits a limited number of "escape" mutations that are associated with resistance to nucleotide analogs such as AZT. However, the vast majority of immunological targets within the protein are preserved even in the drug escape mutants.

Example 54 Construction of HBV Polymerase Plasmid

Experiments reported in the literature indicate that HBV polymerase expression has been achieved in tissue culture cells when both the core and polymerase open reading frames are present in a mRNA molecule. It has also been demonstrated that in this situation, mutation of the core ATG did not influence polymerase expression.

The HBV genome is amplified from a plasmid containing a head-to-tail dimer of the ADW HBV strain. Because expression of polymerase only, and not core is desired, the 5' PCR primer is designed to mutate the precore and core translation initiation codons. In addition, this primer also introduces a mutant DR1 sequence to eliminate the possibility of the generation of a replication-competent HBV genomic RNA. This PCR product is placed into a plasmid containing a kanamycin resistance gene and a pBR322 origin of replication. In addition, this plasmid contains a

- 91 -

cytomegalovirus promoter, a Rous sarcoma virus enhancer, and a SV40 polyadenylation signal. The translation initiation codons for surface antigen and the product of the X coding region are mutated to prevent the expression of the HBs and X gene products.

According to another approach to achieve expression of the HBV polymerase, a PCR product encoding the entire polymerase coding region is amplified and cloned into a vector containing a kanamycin resistance gene and a pBR322 origin of replication. In addition, this plasmid contains a cytomegalovirus promoter, a Rous sarcoma virus enhancer, and a SV40 polyadenylation signal. The 5' PCR primer for this amplification contains a cloning site and spans the translational initiation codon of the polymerase gene. The 3' PCR product contains a restriction site for cloning the insert into the expression vector and also is complementary to the traditional stop codon of the HBV polymerase gene and sequences that flank this stop codon. After ligation of this PCR product into a plasmid containing the kanamycin resistance gene, a pBR322 origin of replication, a cytomegalovirus promoter, a Rous sarcoma virus enhancer, and a SV40 polyadenylation signal, the translation initiation codons for the Hepatitis B surface antigen and X genes are mutated to prevent expression of these gene products. An alternative strategy is used similar to that described above, however, the 3' PCR primer in this case includes the HBVpolyA signal and sequences which flank this signal. This 3' primer is used in the case that sequences including and/or surrounding the HBV polyA signal are important for expression. A mutational analysis has demonstrated that the function of the HBV polymerase gene product can be eliminated by particular nucleotide changes (Radziwell, G. et al. (1990) *J. Virol.* 64(2):613-620). Before utilizing a plasmid constructed as described above, the expressed polymerase can be mutated by the introduction of one of these mutations or others that are analogous.

Example 55

- 92 -

Granulocyte-macrophage colony stimulating factor (GM-CSF) exhibits stimulatory effects on a variety of cell lineages including neutrophils, monocyte/macrophages and eosinophils. The effects of GM-CSF make it an attractive therapeutic model. GM-CSF has been approved by the FDA for use in the autologous bone marrow transplantation and clinical trials have been initiated to test the efficacy in the treatment of various neutropenias. Presently, GM-CSF is administered as a protein which usually requires that it be administered in multiple doses. Proteins must be produced and purified.

An alternative approach to the use of GM-CSF protein is the direct administration of a gene construct which contains a gene encoding GM-CSF in conjunction with the administration of bupivacaine. The genetic construct is constructed by PCR of a GM-CSF gene including signal sequence. The genetic construct preferably contains a kanamycin resistance gene (aminoglycoside 3'-phosphotransferase gene), a bacterial origin of replication, sequences that support expression of the GM-CSF coding region in the cells that the plasmid is introduced into such as the vectors described as backbones in Example 46. The plasmid preferably contains a mammalian origin of replication induced by the cellular replication associated with bupivacaine administration. If the EBV origin of replication is used, the sequence that encodes the nuclear antigen EBNA-1 is also included with the appropriate regulatory sequences. The primers for PCR amplification of the insert contain restriction enzyme sites to allow cloning into the expression vector and are complementary to the 5' and 3' ends of the GM-CSF coding sequences. The PCR reaction is performed with a cDNA clone as described in Lee et al. *Proc. Natl. Acad. Sci., USA* 82:4360-4364.

Example 56

Chronic myelogenous leukemia (CML) is a clonal myeloproliferative disorder of the hematopoietic stem cells associated with the Philadelphia chromosome; a chromosome

- 93 -

abnormality resulting from translocation between chromosomes 9 and 22. The breakpoints on chromosome 22 are clustered in a 6 kb region termed the breakpoint cluster region (BCR), while on chromosome 9, the breakpoints are scattered throughout a 90 kb region upstream from c-abl exon 2. The various 9:22 translocations that result can be subdivided into two types: K28 translocations and L6 translocations. Transcription through the *bcr-abl* translocation results in the generation of fusion mRNAs. Antisense targeted to the *bcr-abl* junction of the mRNAs has been demonstrated to decrease the ability of hematopoietic cells obtained from CML patients to form colonies.

A genetic construction that encodes the antisense is administered together with bupivacaine to the cells of an individual suffering from CML *ex vivo*. The treated cells are then reintroduced into the individual.

Example 57

Gene constructs useful in pharmaceutical kits and compositions for vaccination against and treatment for HBV are constructed with vectors described as backbones in Example 46. The plasmids contain HBV structural genes, particularly genes that encode HBV surface antigen and/or HBV core antigen core and/or HBV precore antigen.

Example 58

Gene constructs useful in pharmaceutical kits and compositions for vaccination against and treatment for HCV are constructed with vectors described as backbones in Example 46. The plasmids contain HCV structural genes, particularly genes that encode HCV core protein and/or HCV envelope protein.

Example 59

The gene construct pREV was designed which contains a nucleotide sequence that encodes HIV rev as the sole target protein. The coding sequence of rev is cloned into Backbone A described in Example 46 from BBG35 (RD Systems Inc. Minneapolis, MN) which contains the coding region of rev from HIV strain HX3B in pUC19.

- 94 -

Tabl 1

Picornavirus Family	
	Genera: Rhinoviruses: (Medical) responsible for ~ 50% cases of the common cold.
5	Etheroviruses: (Medical) includes polioviruses, coxsackieviruses, echoviruses, and human enteroviruses such as hepatitis A virus.
10	Aphoviruses: (Veterinary) these are the foot and mouth disease viruses.
	Target antigens: VP1, VP2, VP3, VP4, VPG
Calcivirus Family	
	Genera: Norwalk Group of Viruses: (Medical) these viruses are an important causative agent of epidemic gastroenteritis.
15	
Togavirus Family	
	Genera: Alphaviruses: (Medical and Veterinary) examples include Senilis viruses, Ross River virus and Eastern & Western Equine encephalitis.
20	Reovirus: (Medical) Rubella virus.
Flariviridae Family	
25	Examples include: (Medical) dengue, yellow fever, Japanese encephalitis, St. Louis encephalitis and tick borne encephalitis viruses.
Hepatitis C Virus: (Medical) these viruses are not placed in a family yet but are believed to be either a togavirus or a flavivirus. Most similarity is with togavirus family.	
30	Coronavirus Family: (Medical and Veterinary)
	Infectious bronchitis virus (poultry)
	Porcine transmissible gastroenteric virus (pig)
35	Porcine hemagglutinating encephalomyelitis virus (pig)
	Feline infectious peritonitis virus (cats)
	Feline enteric coronavirus (cat)
40	Canine coronavirus (dog)
	The human respiratory coronaviruses cause ~40 cases of common cold. EX. 224E, OC43
	Note - coronaviruses may cause non-A, B or C hepatitis
	Target antigens:
45	E1 - also called M or matrix protein
	E2 - also called S or Spike protein
	E3 - also called HE or hemagglutinin-esterase glycoprotein (not present in all coronaviruses)
50	N - nucleocapsid

- 95 -

Rhabdovirus Family

Genera: Vesiliovirus
 Lyssavirus: (medical and veterinary)
 rabies

5 Target antigen: G protein
 N protein

Filoviridae Family: (Medical)

Hemorrhagic fever viruses such as Marburg
 and Ebola virus

10 Paramyxovirus Family:

Genera: Paramyxovirus: (Medical and Veterinary)
 Mumps virus, New Castle disease virus
 (important pathogen in chickens)
 Morbillivirus: (Medical and Veterinary)
 15 Measles, canine distemper
 Pneumovirus: (Medical and Veterinary)
 Respiratory syncytial virus

Orthomyxovirus Family (Medical)

The Influenza virus

20 Bungavirus Family

Genera: Bungavirus: (Medical) California
 encephalitis, LA Crosse
 Phlebovirus: (Medical) Rift Valley Fever
 Hantavirus: Puumala is a hantavirus fever
 25 virus
 Nairovirus (Veterinary) Nairobi sheep
 disease
 Also many unassigned bungaviruses

Arenavirus Family (Medical)

30 LCM, Lassa fever virus

Reovirus Family

Genera: Reovirus: a possible human pathogen
 Rotavirus: acute gastroenteritis in
 children
 35 Orbiviruses: (Medical and Veterinary)
 Colorado Tick fever, Lebombo (humans)
 equine encephalosis, blue tongue

Retrovirus Family**Sub-Family:**

40 Oncorivirinal: (Veterinary) (Medical)
 feline leukemia virus, HTLV I and HTLV II
 Lentivirinal: (Medical and Veterinary)
 HIV, feline immunodeficiency virus,
 equine infections, anemia virus
 45 Spumavirinal

Papovavirus Family**Sub-Family:**

Polyomaviruses: (Medical) BKU and JCU viruses

- 96 -

Sub-Family:

Papill mavirus: (Medical) many viral
types associated with cancers or
malignant progressi n of papilloma

5 Adenovirus (Medical)

EX AD7, ARD., O.B. - cause respiratory disease
- some adenoviruses such as 275 cause
enteritis

10 Parvovirus Family (Veterinary)

Feline parvovirus: causes feline enteritis
Feline panleucopeniavirus
Canine parvovirus
Porcine parvovirus

15 Herpesvirus Family

Sub-Family: alphaherpesviridue

Genera: Simplexvirus (Medical)

HSVI, HSVII

Varicellovirus: (Medical - Veterinary)
pseudorabies - varicella zoster

20 Sub-Family - betaherpesviridue

Genera: Cytomegalovirus (Medical)

HCMV

Muromegalovirus

Sub-Family: Gammaherpesviridue

25 Genera: Lymphocryptovirus (Medical)

EBV - (Burkitts lympho)

Rhadinovirus

Poxvirus Family

Sub-Family: Chordopoxviridue (Medical - Veterinary)

30 Genera: Variola (Smallpox)

Vaccinia (Cowpox)

Parapoxivirus - Veterinary

Auipoxvirus - Veterinary

Capripoxvirus

35 Leporipoxvirus

Suipoxvirus

Sub-Family: Entemopoxviridue

Hepadnavirus Family

Hepatitis B virus

40 Unclassified

Hepatitis delta virus

- 97 -

Table 2

Bacterial pathogens	
5	Pathogenic gram-positive cocci include: pneumococcal; staphylococcal; and streptococcal.
	Pathogenic gram-negative cocci include: meningococcal; and gonococcal.
10	Pathogenic enteric gram-negative bacilli include: enterobacteriaceae; pseudomonas, acinetobacteria and eikenella; melioidosis; salmonella; shigellosis; hemophilus; chancroid; brucellosis; tularemia; yersinia (pasteurella); streptobacillus moniliformis and spirillum; listeria monocytogenes; erysipelothrix rhusiopathiae; diphtheria; cholera; anthrax; donovanosis
	(granuloma inguinale); and bartonellosis.
20	Pathogenic anaerobic bacteria include: tetanus; botulism; other clostridia; tuberculosis; leprosy; and other mycobacteria. Pathogenic spirochetal diseases include: syphilis; treponematoses: yaws, pinta and endemic syphilis; and leptospirosis.
	Other infections caused by higher pathogen bacteria and pathogenic fungi include: actinomycosis; nocardiosis; cryptococcosis, blastomycosis, histoplasmosis and coccidioidomycosis; candidiasis, aspergillosis, and mucormycosis; sporotrichosis; paracoccidioidomycosis, petriellidiosis, torulopsosis, mycetoma and chromomycosis; and dermatophytosis.
30	Rickettsial infections include rickettsial and rickettsioses.
35	Examples of mycoplasma and chlamydial infections include: mycoplasma pneumoniae; lymphogranuloma venereum; psittacosis; and perinatal chlamydial infections.
	Pathogenic eukaryotes
40	Pathogenic protozoans and helminths and infections thereby include: amebiasis; malaria; leishmaniasis; trypanosomiasis; toxoplasmosis; pneumocystis carinii; babesiosis; giardiasis; trichinosis; filariasis; schistosomiasis; nematodes; trematodes or flukes; and cestode (tapeworm) infections.

- 98 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Weiner, David B.
Williams, William V.
Wang, Bin
Coney, Leslie R.
Merva, Michael J.
Zurawski, Vincent R., Jr.
- 10 (ii) TITLE OF INVENTION: Compositions and Methods for Delivery of
Genetic Material
- (iii) NUMBER OF SEQUENCES: 52
- 15 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Woodcock Washburn Kurtz Mackiewicz & Norris
(B) STREET: One Liberty Place 46th Floor
(C) CITY: Philadelphia
(D) STATE: Pennsylvania
(E) COUNTRY: USA
(F) ZIP: 19103
- 20 (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25 mb-MD/JAF
- 25 (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:
- 30 (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/125,012
(B) FILING DATE: 21-SEP-1993
(C) CLASSIFICATION:
- 35 (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/124,962
(B) FILING DATE: 21-SEP-1993
(C) CLASSIFICATION:
- 40 (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/093,235
(B) FILING DATE: 15-JUL-1993
(C) CLASSIFICATION:
- 45 (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/029,336
(B) FILING DATE: 11-MAR-1993
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/008,342
(B) FILING DATE: 26-JAN-1993
(C) CLASSIFICATION:
- 50 (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: DeLuca, Mark
(B) REGISTRATION NUMBER: 33,229
(C) REFERENCE/DOCKET NUMBER: APOL-0013
- 55 (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 215-568-3100
(B) TELEFAX: 215-568-3429

- 99 -

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AGGCGTCTCG AGACAGAGGA GAGCAAGAAA TG

32

10 (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TTTCCCTCTA GATAAGCCAT CCAATCACAC

30

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAAGGATCCA TGAAAAATA TTTATTGGG

29

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ACTGTCGACT TATTTTAAAG CGTTTTTAAG

30

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 100 -

- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
- GCCAGTTTTG GATCCTTAAA AAAGGCTTGG 30
- (2) INFORMATION FOR SEQ ID NO:6:
- 5 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
- TTGTGAGGGA CAGAATTCCA ATCAGGG 27
- (2) INFORMATION FOR SEQ ID NO:7:
- 15 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
- CAGTGATATC CCGGGAGACT CCTC 24
- (2) INFORMATION FOR SEQ ID NO:8:
- 25 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:
- 30 GAATAGAAGA ACTCCTCTAG AATTC 25
- (2) INFORMATION FOR SEQ ID NO:9:
- 35 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:
- GCCTTAGGCG GATCCTATGG CAGGAAG 27

- 101 -

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

TAAGATGGGT GGCCATGGTG AATT

24

10 (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Cys Asn Thr Arg Lys Arg Ile Arg Ile Gln Arg Gly Pro Gly Arg Ala
 1 5 10 15

20 Phe Val Thr Ile Gly Lys
 20

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Tyr Asn Lys Arg Lys Arg Ile His Ile Gln Arg Gly Pro Gly Arg Ala
 1 5 10 15

30 Phe Tyr Thr Thr Lys Asn Ile Ile Cys
 20 25

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

40 Cys Arg Ile Lys Gln Phe Ile Asn Met Trp Gln Glu Val Gly Lys Ala
 1 5 10 15

- 102 -

Met Thr Ala Pro Pro Ile Ser Gly Ile Arg Cys
20 25

(2) INFORMATION FOR SEQ ID NO:14:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

10 Arg Ile Leu Ala Val Glu Arg Tyr Ile Lys Asp Gln Gln Leu Leu Gly Ile
1 5 10 15
Trp Gly Cys Ser Gly Lys Leu Ile Cys
20 25

(2) INFORMATION FOR SEQ ID NO:15:

- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TTGTTTAACT TTTGATCGAT CCATTCC

27

(2) INFORMATION FOR SEQ ID NO:16:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GATTTGTATC GATGATCTGA C

21

(2) INFORMATION FOR SEQ ID NO:17:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

- 103 -

TG TAGTAGCA AAAGAAATAG TTAAG

25

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

10 AATTCTTAAC TATTCTTTT GCTAC

25

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

ATTGTGCGAC TGGTTTCAGC CTGCCATGGC AGGAAGAACC

40

20 (2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ACGACGCGTA TTCTTTAGCT CCTGACTCC

29

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

35

- 104 -

GCTGACGGTA GCGGCCGCAC AATT

24

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

10 GTATTAAGCG GCCGCAATTG TT

22

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 78 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

AAAAAGCTTC GCGGATCCGC GTTGCGGCCG CAACCGGTCA CCGGCGACGC GTCGGTCGAC

60

20 CGGTCATGGC TGGGCCCC

78

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CCCAAGCTTA GACATGATAA GATACATTG

29

30 (2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

- 105 -

CTAGCAGCTG GATCCCAGCT TC

22

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

10 GGATTCTGG GGATCCAAGC TAGT

24

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

TATAGGATCC GCGCAATGAA AGACCCACCC T

31

20 (2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

ATATGGATCC GCAATGAAAG ACCCCGCTG A

31

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

TAAAGCGGCC GTCCTATGG CAGGAAGACC

30

(2) INFORMATION FOR SEQ ID NO:30:

- 106 -

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

ATTACGCGTC TTATGCTTCT AGCCAGGCAC AATG

34

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ATTACGCGTT TATTACAGAA TGGAAAACAG ATGGCAGGTG

40

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

25

ATTACGCGTT ATTGCAGAAT TCTTATTATG GC

32

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

35 GAGGCTTGGA GAGGATTATA GAAGTACTGC AAGAGCTG

38

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid

- 107 -

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

5 GAATCCTCTC CAAGCCTCAG CTA CTGCTAT AGCTGTGGC 39

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

AAAAATAAAG CGGCCGCTCC TATGGCAGGA AGAGAAGCG 39

15 (2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

AAAAAATTAC GCGTCTTATG CTTCTAGCCA GGCACAATG 39

(2) INFORMATION FOR SEQ ID NO:37:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

CCCAAGCTTG GGAATGCTCT GCCAGTGTTA C 31

(2) INFORMATION FOR SEQ ID NO:38:

35 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

- 108 -

GGGGGCCCGGA AGGGCACAAAT AAAACTGTCT GCTTAC

36

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

10 CCTGATTGAG GTGAAAATAT TGTTGATGCG CTG

33

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 111 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

AACATCAATA CAACCTATTA ATTTCCCTC GTCAAAAATA AGGTTATCAA GTGAGAAATC

60

20 ACCATCAGTG ACGACTGAAT CCGGTGAGAA TGGCAAAAGT TTATGCATTT C

111

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 55 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

CTAGCGCGGG GATCCGCGTT GCGGCGCAA AAAGTCGACG GCGGACGCGT AAAAA

55

30 (2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 55 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GATCTTTTTA CGCGTCGCCC GTCGACTTTT TCGGCGCGCA ACGCGGATCC CCGCG

55

(2) INFORMATION FOR SEQ ID NO:43:

- 109 -

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 48 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 5 (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:
ATGTCGACTG GTTTCAGCCT GCCATGGCAG GAAGAAGC 48
- (2) INFORMATION FOR SEQ ID NO:44:
- 10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 15 (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:
CCCCACGACG CGTCTATTCT TTAGCTCCTG ACTCC 35
- (2) INFORMATION FOR SEQ ID NO:45:
- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:
TTTGCCGCCG CGTAAGTGA GAGAGATGCT GCGAG 35
- (2) INFORMATION FOR SEQ ID NO:46:
- 30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:
35 CTGGTGGGGC TGTGGCTCT G 21
- (2) INFORMATION FOR SEQ ID NO:47:
- 40 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 80 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA

- 110 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

AATTTAATAA GTAAGTAAGT GTCATATGTT TGTTTGAATT CTGCAACAAC TGCTGTTTAT 60
CCATTTTCAG AATTGGGTG 80

(2) INFORMATION FOR SEQ ID NO:48:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 80 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

TCGACACCCA ATTCTGAAAA TGGATAAACA GCACTTGTTG CAGAATTCAA ACAAACATAT 60
GACACTTACT TACTTATTA 80

(2) INFORMATION FOR SEQ ID NO:49:

- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

GGGGTTTTTG GGCATATGTA TGAGGGACAA TTGAGAAGT G 41

(2) INFORMATION FOR SEQ ID NO:50:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

AAGCTTGTGG AATTCTTAAT TTCTCTGTCC GGGGTTTTTG GGCATATGTA TGAGGGACAT 60
TCGAGAAGTG 70

(2) INFORMATION FOR SEQ ID NO:51:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 111 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CAGTATCTGG CATGGGTAC

29

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

CCATGCCAGA TACTGGTAC

29

- 112 -

Claims

1. A method of introducing genetic material into cells of an individual comprising the steps of:
 - a) contacting cells of said individual with a polynucleotide function enhancer;
 - b) administering to cells of said individual, a nucleic acid molecule;wherein said nucleic acid molecule is free of retroviral particles.
2. The method of claim 1 wherein said polynucleotide function enhancer is a bupivacaine.
3. The method of claim 1 wherein said nucleic acid molecule comprises a nucleotide sequence that encodes a protein and is operably linked to regulatory sequences.
4. The method of claim 1 wherein said nucleic acid molecule comprises a nucleotide sequence that encodes a protein which comprises at least one epitope that is identical or substantially similar to an epitope of an antigen against which an immune response is desired, said nucleotide sequence operably linked to regulatory sequences.
5. A method of immunizing an individual against a pathogen comprising the steps of:
 - a) contacting cells of said individual with a polynucleotide function enhancer;
 - b) administering to cells of said individual, a nucleic acid molecule that comprises a nucleotide sequence that encodes a protein which comprises at least one epitope that is identical or substantially similar to an epitope of a pathogen antigen, said nucleotide sequence being operably linked to regulatory sequences;wherein said nucleic acid molecule is free of retroviral particles and said nucleotide sequence is capable of being expressed in said cells.

- 113 -

6. The method of Claim 5 where in said polynucleotide functi n enhancer is bupivacaine.
7. The method of Claim 5 wherein said nucleic acid molecule is a DNA molecule.
- 5 8. The method of Claim 5 wherein said protein is a pathogen antigen or a fragment thereof.
9. The method of Claim 5 wherein said nucleic acid molecule is administered intramuscularly.
10. The method of Claim 5 wherein said pathogen is a
10 virus selected from the group consisting of: human immunodeficiency virus, HIV; human T cell leukemia virus, HTLV; influenza virus; hepatitis A virus, HAV; hepatitis B virus, HBV; hepatitis C virus, HCV; human papilloma virus, HPV; Herpes simplex 1 virus, HSV1; Herpes simplex 2 virus,
15 HSV2; Cytomegalovirus, CMV; Epstein-Barr virus, EBV; rhinovirus; and, coronavirus.
11. The method of Claim 5 wherein said pathogen is HIV and said nucleic acid molecule comprises a nucleotide sequence sequences that encodes an HIV protein.
- 20 12. The method of Claim 5 wherein said pathogen is HIV and said nucleic acid molecule comprises a nucleotide sequence sequences that encodes more than one HIV structural protein.
13. The method of Claim 5 wherein said pathogen is HIV and said nucleic acid molecule comprises a nucleotide sequence
25 sequences that encodes more than one HIV regulatory protein.
14. The method of Claim 5 wherein at least two or more different nucleic acid molecules are administered to different cells of an individual; said different nucleic acid molecules

- 114 -

each comprise nucleotide sequences encoding one or more pathogen antigens of the same pathogen.

15. The method of Claim 5 wherein said polynucleotide function enhancer and said nucleic acid molecule are administered simultaneously.

16. The method of Claim 5 wherein:
said individual is a human;
said polynucleotide function enhancer is bupivacaine;

10 said pathogen is human immunodeficiency virus;
said nucleic acid molecule is DNA and comprises a DNA sequence that encodes HIV structural proteins gag and pol with a deletion of the psi, said DNA sequence that encodes gag and pol operably linked to a rous sarcoma virus enhancer,
15 a cytomegalovirus immediate early promoter and an SV40 minor polyadenylation signal and optionally an SV40 origin of replication.

17. The method of Claim 16 wherein said DNA sequence further comprises an HIV rev response element and a deletion
20 of HIV integrase.

18. The method of Claim 17 wherein said DNA sequence further comprises an HIV splice acceptor.

19. The method of Claim 16 wherein said DNA molecule further comprises a DNA sequence that encodes sequence further
25 comprises HIV rev operably linked to an SV40 promoter and an SV40 minor polyadenylation signal and optionally an SV40 origin of replication.

20. The method of Claim 19 wherein said DNA sequence that encodes rev additionally encodes HIV vpu and HIV env.

- 115 -

21. The method of Claim 19 wherein said DNA sequence that encodes *gag* and *pol* further comprises an HIV rev response element and a deletion of HIV integrase.

22. The method of Claim 21 wherein said DNA sequence
5 that encodes *gag* and *pol* further comprises an HIV splice acceptor.

23. The method of Claim 5 wherein:
said individual is a human;
said polynucleotide function enhancer is
10 bupivacaine;
said pathogen is human immunodeficiency virus;
said nucleic acid molecules is DNA and
comprises a DNA sequence that encodes HIV proteins *rev*, *vpu*
and *env* operably linked to a rous sarcoma virus enhancer, a
15 cytomegalovirus immediate early promoter and an SV40 minor
polyadenylation signal and optionally an SV40 origin of
replication.

24. The method of Claim 5 wherein:
said individual is a human;
20 said polynucleotide function enhancer is
bupivacaine;
said pathogen is human immunodeficiency virus;
two different DNA molecules are administered
to different cells of an individual;
25 one of said nucleic acid molecules is DNA and
comprises a DNA sequence that encodes HIV structural proteins
gag and *pol* with an deletion of the *psi*, said DNA sequence
that encodes *gag* and *pol* operably linked to a rous sarcoma
virus enhancer, a cytomegalovirus immediate early promoter and
30 an SV40 minor polyadenylation signal and optionally an SV40
origin of replication; and
the other of said nucleic acid molecules is DNA
and comprises a DNA sequence that encodes HIV proteins *rev*,
vpu and *env* operably linked to a rous sarcoma virus enhancer,

- 116 -

a cytomegalovirus immediate early promoter and an SV40 minor polyadenylation signal and optionally an SV40 origin of replication.

25. A method of immunizing a human against HIV
5 comprising the steps of:

administering to cells of said human, two different nucleic acid molecules; each of said nucleic acid molecules comprising a nucleotide sequence that encodes a protein which comprises at least one epitope that is identical or
10 substantially similar to an epitope of at least one HIV antigen operatively linked to regulatory sequences; said nucleotide sequences being capable of being expressed in said cells; nucleotide sequences of each of said different nucleic acid molecules encode different proteins; said proteins
15 comprise at least one epitope that is identical or substantially similar to an epitope of at least one of the HIV proteins encoded by HIV genes selected from the group consisting of *gag*, *pol* and *env*.

26. A method of immunizing an individual against a
20 disease comprising the steps of:

a) contacting cells of said individual with a polynucleotide unction enhancer;

b) administering to cells of said individual, a nucleic acid molecule comprising a nucleotide sequence that
25 encodes a target protein which comprises an epitope identical or substantially similar to an epitope of a protein associated with cells that characterize said disease operatively linked to regulatory sequences;

wherein said nucleic acid molecule is free of
30 retroviral particles and capable of being expressed in said cells.

27. The method of Claim 26 wherein said polynucleotide function enhancer is bupivacaine.

- 117 -

28. The method of Claim 26 wherein said disease is characterized by hyperproliferating cells.

29. The method of Claim 26 wherein said disease is an autoimmune disease.

5 30. The method of Claim 26 wherein said nucleic acid molecule is a DNA molecule.

31. The method of Claim 26 wherein said nucleic acid molecule is administered intramuscularly.

32. The method of Claim 26 wherein said nucleic acid
10 molecule comprises a nucleotide sequence that encodes a target protein selected from the group consisting of: protein products of oncogenes *myb*, *myc*, *fyn*, *ras*, *sarc*, *neu* and *trk*; protein products of translocation gene *bcl/abl*; P53; EGRF; variable regions of antibodies made by B cell lymphomas; and
15 variable regions of T cell receptors of T cell lymphomas.

33. The method of Claim 26 wherein said protein is selected from the group consisting of: variable regions of antibodies involved in B cell mediated autoimmune disease; and variable regions of T cell receptors involved in T cell
20 mediated autoimmune disease.

34. An pharmaceutical immunizing kit comprising:

a) a first inoculant comprising:

i) a pharmaceutically acceptable carrier

or diluent; and,

25 ii) a first nucleic acid molecule comprising a nucleotide sequence that encodes at least one HIV protein operatively linked to regulatory sequences; wherein said nucleotide sequence is capable of being expressed in human cells;

30 b) a second inoculant comprising:

- 118 -

i) a pharmaceutically acceptable carrier or diluent; and,

ii) a second nucleic acid molecule comprising a nucleotide sequence that encodes at least one HIV protein operatively linked to regulatory sequences; wherein
5 said nucleotide sequence is capable of being expressed in human cells;

wherein said first nucleic acid molecule is not identical to said second nucleic acid molecule.

10 35. The pharmaceutical kit of Claim 34 further comprising:

c) a third inoculant comprising bupivacaine.

36. The pharmaceutical composition of claim 34 wherein taken together said first nucleic acid molecule and said
15 second nucleic acid molecule encode HIV proteins *gag*, *pol* and *env*.

37. A pharmaceutical composition comprising

a) a DNA molecule that comprises a DNA sequence that encodes HIV structural proteins *gag* and *pol* with a
20 deletion of the *psi*, said DNA sequence that encodes *gag* and *pol* operably linked to a rous sarcoma virus enhancer, a cytomegalovirus immediate early promoter and an SV40 minor polyadenylation signal and optionally an SV40 origin of replication; and

25 b) a polynucleotide function enhancer.

38. The pharmaceutical composition of claim 37 wherein said cell stimulating compound is bupivacaine

39. The pharmaceutical composition of claim 37 wherein said DNA sequence further comprises an HIV rev response
30 element and a deletion of HIV integrase.

- 119 -

40. The pharmaceutical composition of claim 37 wherein said DNA sequence further comprises an HIV splice acceptor.

41. The pharmaceutical composition of claim 37 wherein said DNA molecule further comprises a DNA sequence that
5 encodes sequence further comprises HIV rev operably linked to an SV40 promoter and an SV40 minor polyadenylation signal and optionally an SV40 origin of replication.

42. The pharmaceutical composition of claim 41 wherein said DNA sequence that encodes rev additionally encodes HIV
10 vpu and HIV env.

43. The pharmaceutical composition of claim 42 wherein said DNA sequence that encodes gag and pol further comprises an HIV rev response element and a deletion of HIV integrase.

44. The method of Claim 42 wherein said DNA sequence
15 that encodes gag and pol further comprises an HIV splice acceptor.

45. A pharmaceutical composition that comprises
a) a DNA molecule that comprises a DNA sequence that encodes HIV proteins rev, vpu and env operably linked to
20 a rous sarcoma virus enhancer, a cytomegalovirus immediate early promoter and an SV40 minor polyadenylation signal and optionally an SV40 origin of replication; and
b) a polynucleotide function enhancer.

46. The pharmaceutical composition of claim 45 wherein
25 said polynucleotide function enhancer is bupivacaine

47. An pharmaceutical immunizing kit comprising:
a) a first inoculant comprising:
i) a first pharmaceutical composition comprising a DNA molecule that comprises a DNA sequence that
30 encodes HIV structural proteins gag and pol with an deletion

- 120 -

of the *psi*, said DNA sequence that encodes *gag* and *pol* operably linked to a rous sarcoma virus enhancer, a cytomegalovirus immediate early promoter and an SV40 minor polyadenylation signal and optionally an SV40 origin of replication; and

ii) a polynucleotide function enhancer;

and

b) a second inoculant comprising:

i) a second pharmaceutical composition comprising a DNA molecule that comprises a DNA sequence that encodes HIV proteins *rev*, *vpu* and *env* operably linked to a rous sarcoma virus enhancer, a cytomegalovirus immediate early promoter and an SV40 minor polyadenylation signal and optionally an SV40 origin of replication; and

ii) a polynucleotide function enhancer.

48. The pharmaceutical immunizing kit of claim 47 wherein said polynucleotide function enhancer is bupivacaine.

49. A method of treating an individual suspected of suffering from a disease comprising the steps of:

a) contacting cells of said individual with a polynucleotide function enhancer;

b) administering to cells of said individual, a nucleic acid molecule comprising a nucleotide sequence that encodes a protein whose presence will compensate for a missing, non-functional or partially functioning protein or produce a therapeutic effect on the individual, said nucleotide sequence operatively linked to regulatory sequences;

wherein said nucleic acid molecule is free of retroviral particles and capable of being expressed in said cells.

50. The method of Claim 49 wherein said polynucleotide function enhancer is bupivacaine.

- 121 -

51. The method of Claim 49 wherein said nucleic acid molecule is a DNA molecule.

52. The method of Claim 49 wherein said nucleic acid molecule is administered intramuscularly.

5 53. The method of Claim 49 wherein said nucleic acid molecule comprises a nucleotide sequence that encodes a protein selected from the group consisting of: enzymes, structural proteins, cytokines, lymphokines and growth factors.

10 54. A pharmaceutical composition comprising:
i) a nucleic acid molecule that comprises a nucleotide sequence which encodes a protein selected from the group consisting of: proteins which comprises at least one epitope that is identical or substantially similar to an
15 epitope of a pathogen antigen; proteins which comprises an epitope identical or substantially similar to an epitope of a protein associated with hyperproliferating cells; proteins which comprises an epitope identical or substantially similar to an epitope of a protein associated with cells that
20 characterize an autoimmune disease; proteins whose presence will compensate for a missing, non-functional or partially functioning protein in an individual; and proteins that produce a therapeutic effect on an individual; and
ii) a polynucleotide function enhancer;
25 wherein said pharmaceutical composition is free of retroviral particles.

55. The method of Claim 54 wherein said polynucleotide function enhancer is bupivacaine.

56. A pharmaceutical kit comprising:
30 i) a container that comprises a nucleic acid molecule that comprises a nucleotide sequence which encodes a protein selected from the group consisting of: proteins

- 122 -

which comprises at least one epitope that is identical or substantially similar to an epitope of a pathogen antigen; proteins which comprises an epitope identical or substantially similar to an epitope of a protein associated with hyperproliferating cells; proteins which comprises an epitope identical or substantially similar to an epitope of a protein associated with cells that characterize an autoimmune disease; proteins whose presence will compensate for a missing, non-functional or partially functioning protein in an individual; and proteins that produce a therapeutic effect on an individual; and

ii) a container that comprises a polynucleotide function enhancer; wherein said pharmaceutical kit is free of retroviral particles.

57. The method of Claim 56 wherein said polynucleotide function enhancer is bupivacaine.

1/10




FIG. 1A

2 / 10




FIG. 1B




FIG. 2

3 / 10




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E

SUBSTITUTE SHEET (RULE 26)

4 / 10



FIG. 4A




FIG. 4B




FIG. 4C



FIG. 4D




FIG. 4E



FIG. 4F




FIG. 4G

6/10




FIG. 5

7/10




FIG. 6




FIG. 7

SUBSTITUTE SHEET (RULE 26)

8/10




FIG. 8A




FIG. 8B

SUBSTITUTE SHEET (RULE 26)




FIG. 8C




FIG. 8D

SUBSTITUTE SHEET (RULE 26)

10 / 10




FIG. 9

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/00899

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : A61K 48/00; C12N 15/12, 15/31, 15/33, 15/48

US CL : 435/69.3, 320.1; 424/93R, 93A, 93B; 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.3, 320.1; 424/93R, 93A, 93B; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, BIOSIS, EMBASE, DERWENT

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	WO, A, 90/11092 (FELGNER et al.) 04 October 1990, see entire document.	1, 3 - 5, 7 - 11, 15, 26, 28-31 ----- 2, 6, 12-14, 16- 25, 27, 32-57
Y	Nature, Volume 352, issued 29 August 1991, G. Acsadi et al., "Human Dystrophin Expression in mdx Mice after Intramuscular Injection of DNA Constructs", pages 815-818, see entire document.	1-57



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

26 APRIL 1994

Date of mailing of the international search report

09 MAY 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Michael S. Tuscan, Ph.D.

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US94/00899

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	American Journal of Physiology, Volume 258, Number 3, issued March 1990, D.B. Thomason et al., "Stable Incorporation of a Bacterial Gene in Adult Rat Skeletal Muscle In Vivo", pages C578-C581, see entire document.	1-57
Y	Nature, Volume 349, issued 24 January 1991, P.L. Felgner et al., "Gene Therapeutics", pages 351-352, see entire document.	1-57
Y	Science, Volume 247, issued 23 March 1990, J.A. Wolf et al, "Direct Gene Transfer into Mouse Muscle In Vivo", pages 1465-1467, see entire document.	1-57
Y	Journal of Virology, Volume 64, Number 5, issued May 1990, A. Aldovini et al., "Mutations of RNA and Protein Sequences Involved in Human Immunodeficiency Virus Type 1 Packaging Result in Production of Noninfectious Virus", pages 1920-1926, see entire document.	24, 37-44, 47-48
Y	Biochemical and Biophysical Research Communications, Volume 185, Number 3, issued 30 June 1992, M. Schauer et al., "The N-Terminal Region of HIV-1 Integrase is Required For Integration Activity, But Not for DNA Binding", pages 874-880, see entire document.	17-18,21-22,39,43
Y	AIDS Research and Human Retroviruses, Volume 8, Number 3, issued March 1992, R.C. Desrosiers et al., "HIV with Multiple Gene Deletions as a Live Attenuated Vaccine for AIDS", pages 411-421, see entire document.	1-57
Y	Nucleic Acids Research, Volume 20, Number 13, issued 11 July 1992, D. Ronen et al, "Expression of Wild-Type and Mutant p53 Proteins by Recombinant Vaccinia Viruses", pages 3435-3441, see entire document.	28, 31-32